

MATRIX UNITS AND GENERIC DEGREES FOR THE ARIKI-KOIKE ALGEBRAS

ANDREW MATHAS

ABSTRACT. We compute the generic degrees of the Ariki–Koike algebras by first constructing a basis of matrix units in the semisimple case. As a consequence, we also obtain an explicit isomorphism from any semisimple Ariki–Koike algebra to the group algebra of the corresponding complex reflection group.

1. INTRODUCTION

The cyclotomic Hecke algebras were introduced by Ariki and Koike [2, 4] and Broué and Malle [7]. It is conjectured [7] that these algebras play a rôle in the representation theory of reductive groups similar to (but more complicated than) that played by the Iwahori–Hecke algebras (see, for example, [8]). In particular, it should be possible to use these algebras to compute the degrees (and more generally characters) of certain representations of reductive groups; more precisely, we can attach a polynomial to each irreducible representation $\mathcal{H}$, called its generic degree, and appropriate specialisations of this polynomial should compute the dimensions of the corresponding irreducible representations of the finite groups of Lie type.

The purpose of this paper is to compute the generic degrees for the cyclotomic Hecke algebras of type $G(r, 1, n)$; these polynomials have also been computed by Geck, Iancu and Malle [15]. Further results of Malle [19] and Malle and the author [21] give the generic degrees for all of the cyclotomic Hecke algebras corresponding to imprimitive complex reflection groups. Malle [20] has recently computed the generic degrees for cyclotomic algebras for the primitive complex reflection groups (modulo the assumption that the corresponding cyclotomic Hecke algebras are symmetric), so this completes the calculation of the generic degrees of the cyclotomic Hecke algebras associated with complex reflection groups.

Two important special cases of the Ariki–Koike algebras are the Iwahori–Hecke algebras of types A and B ; the generic degrees of these algebras are well-known and were first computed by Hoefsmit [16]. Later Murphy [25] gave an easier derivation of Hoefsmit’s formulae for the generic degrees of the Iwahori–Hecke algebras of type A using different, but related, techniques.

This article is largely inspired by Murphy’s paper [25]; however, with hindsight we are able to take quite a few shortcuts. Along the way we give a new and quite elegant treatment of the representation theory of the semisimple Ariki–Koike algebras. In particular, we explicitly construct the primitive idempotents and the matrix units in the Wedderburn decomposition of $\mathcal{H}$. One of the nice features of our approach is that we use the modular theory (more accurately, the cellular

1991 *Mathematics Subject Classification.* 20C08, 20G05, 33D80.

Key words and phrases. Ariki–Koike algebras, generic degrees and complex reflection groups.

theory) to understand the semisimple case. Another advantage of our approach is that our calculation of the generic degrees is almost entirely algebraic and it is only near of the end of the paper that we need to introduce combinatorial arguments (however, this is not to say that our approach is painless; in addition, showing that our result agrees with [15] is a long and unpleasant calculation).

2. THE ARIKI-KOIKE ALGEBRAS

Let R be a commutative domain with 1 and fix elements $q, Q_1, \dots, Q_r$ in R with q invertible. Let $\mathbf{q} = (q; Q_1, \dots, Q_r)$. The Ariki-Koike algebra $\mathcal{H} = \mathcal{H}_{\mathbf{q}}(n)$ is the unital associative algebra with generators $T_0, T_1, \dots, T_{n-1}$ and relations

$$\begin{aligned} (T_0 - Q_1) \cdots (T_0 - Q_r) &= 0, \\ T_0 T_1 T_0 T_1 &= T_1 T_0 T_1 T_0, \\ (T_i + q)(T_i - 1) &= 0, & \text{for } 1 \leq i \leq n-1, \\ T_{i+1} T_i T_{i+1} &= T_i T_{i+1} T_i, & \text{for } 1 \leq i \leq n-2, \\ T_i T_j &= T_j T_i, & \text{for } 0 \leq i < j-1 \leq n-2. \end{aligned}$$

The Ariki-Koike algebra is a deformation of the group algebra of $W_{r,n}$, where $W_{r,n} = (\mathbb{Z}/r\mathbb{Z}) \wr \mathfrak{S}_n$ is the wreath product of the cyclic group of order r and the symmetric group $\mathfrak{S}_n$ of degree n .

Define elements $L_m = q^{1-m} T_{m-1} \cdots T_1 T_0 T_1 \cdots T_{m-1}$ for $m = 1, 2, \dots, n$; these are analogues of the q -Murphy operators of the Iwahori-Hecke algebras of type **A** [10, 25]. An easy calculation using the relations in $\mathcal{H}$ (see [4, 3.3] and [10, (2.1), (2.2)]) shows that we have the following results.

2.1 *Suppose that $1 \leq i \leq n-1$ and $1 \leq m \leq n$. Then*

- (i) L_i and L_m commute.
- (ii) T_i and L_m commute if $i \neq m-1, m$.
- (iii) T_i commutes with $L_i L_{i+1}$ and $L_i + L_{i+1}$.
- (iv) If $a \in R$ and $i \neq m$ then T_i commutes with $(L_1 - a)(L_2 - a) \cdots (L_m - a)$.

Using the elements T_w and L_m defined above, Ariki and Koike proved the following result which gives a basis for $\mathcal{H}$.

2.2 (Ariki-Koike [4, (3.10)]) *The algebra $\mathcal{H}$ is free as an R -module with basis*

$$\{ L_1^{c_1} L_2^{c_2} \cdots L_n^{c_n} T_w \mid w \in \mathfrak{S}_n \text{ and } 0 \leq c_m \leq r-1 \text{ for } m = 1, 2, \dots, n \}.$$

In particular, $\mathcal{H}$ is free of rank $r^n n!$

We call this basis the Ariki-Koike basis of $\mathcal{H}$. Let $\mathcal{H}(\mathfrak{S}_n)$ be the subalgebra of $\mathcal{H}$ generated by $T_1, \dots, T_{n-1}$. It follows from (2.2) that $\mathcal{H}(\mathfrak{S}_n)$ is isomorphic to the Iwahori-Hecke algebra of $\mathfrak{S}_n$ and is free as an R -module with basis $\{ T_w \mid w \in \mathfrak{S}_n \}$.

Let $\tau: \mathcal{H} \rightarrow R$ be the R -linear map determined by

$$\tau(L_1^{c_1} L_2^{c_2} \cdots L_n^{c_n} T_w) = \begin{cases} 1, & \text{if } c_1 = \cdots = c_n = 0 \text{ and } w = 1, \\ 0, & \text{otherwise,} \end{cases}$$

where $w \in \mathfrak{S}_n$ and $0 \leq c_i < r$ for $i = 1, \dots, n$.

The function τ was introduced by Bremke and Malle [5] who showed that τ is a trace form and that τ is essentially independent on the choice of basis of $\mathcal{H}$. It is not obvious from the definition above that τ coincides with the form introduced by Bremke and Malle; however, this was proved by Malle and the author in [21] where we also showed that τ is non-degenerate whenever $Q_1, \dots, Q_r$ are invertible in R .

For future reference we note the following two important properties of τ ; the first is Bremke and Malle's result that τ is a trace form and the second follows easily from the definition and well-known properties of the trace form τ in the case $r = 1$ (see, for example, [22, Prop. 1.16]).

- 2.3** (i) Suppose $h_1, h_2 \in \mathcal{H}$. Then $\tau(h_1 h_2) = (h_2 h_1)$.
(ii) Suppose that $x, y \in \mathfrak{S}_n$ and that $0 \leq c_i < r$ for $i = 1, \dots, n$. Then

$$\tau(L_1^{c_1} L_2^{c_2} \dots L_n^{c_n} T_x T_y) = \begin{cases} q^{\ell(x)}, & \text{if } c_1 = \dots = c_n = 0 \text{ and } x = y^{-1}, \\ 0, & \text{otherwise.} \end{cases}$$

In this paper we will mainly be concerned with the semisimple Ariki–Koike algebras; these were classified by Ariki [1] who showed that when R is a field $\mathcal{H}$ is semisimple if and only if

$$(2.4) \quad P_{\mathcal{H}}(\mathbf{q}) = \prod_{i=1}^n (1 + q + \dots + q^{i-1}) \cdot \prod_{1 \leq i < j \leq r} \prod_{-n < d < n} (q^d Q_i - Q_j)$$

is a non-zero element of R . For most of what we do it will be enough to assume that R is a ring in which $P_{\mathcal{H}}(\mathbf{q})$ is invertible.

A **multipartition** of n is a ordered r -tuple $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ of partitions $\lambda^{(s)}$ such that $n = \sum_{s=1}^r |\lambda^{(s)}|$; we write $\lambda \vdash n$. In the semisimple case Ariki and Koike [4] constructed an irreducible $\mathcal{H}$ -module S^λ , called a **Specht module**, for each multipartition λ of n . Further, they showed that $\{S^\lambda \mid \lambda \vdash n\}$ is a complete set of pairwise non-isomorphic irreducible $\mathcal{H}$ -modules. Let χ^λ be the character of S^λ .

Assume that $\mathcal{H}$ is semisimple. Then τ can be written as a linear combination of the irreducible characters of $\mathcal{H}$ because τ is a trace form.

2.5. Definition. Suppose that R is a field and that $P_{\mathcal{H}}(\mathbf{q}) \neq 0$. The Schur elements of $\mathcal{H}$ are the elements $s_\lambda(\mathbf{q}) \in R$ such that

$$\tau = \sum_{\lambda} \frac{1}{s_\lambda(\mathbf{q})} \chi^\lambda,$$

where λ runs over the multipartitions of n .

The rational functions $\frac{1}{s_\lambda(\mathbf{q})}$ are also called the weights of $\mathcal{H}$.

The **generic degrees** of $W_{r,n}$ are certain “spetsial” specializations of the rational functions $s_\eta(\mathbf{q})/s_\lambda(\mathbf{q})$, where $\eta = ((n), (0), \dots, (0))$ — so $s_\eta(\mathbf{q})$ is the Schur element corresponding to the trivial representation of $\mathcal{H}$ ($s_\eta(\mathbf{q}) = s_{\eta_1}(\mathbf{q})$ is computed in Example 2.7). The spetsial specializations of the rational functions $s_\eta(\mathbf{q})/s_\lambda(\mathbf{q})$ are polynomials in q with rational coefficients; moreover, for these specializations $s_\eta(\mathbf{q})$ is equal to the Poincaré polynomial of the coinvariant algebra of the reflection representation of $W_{r,n}$. These results are due to Malle and can be found in [19, 20].

In the special case when $r = 1, 2$ the group $W_{r,n}$ is a Weyl group (rather than just a complex reflection group), and here the generic degrees were first computed by Hoefsmit [16]; they can be found, for example, in [8].

One of the motivations for writing this paper was to compute the Schur elements and hence the generic degrees. Ostensibly the Schur elements depend in a non-uniform way upon the choice of $q, Q_1, \dots, Q_r$; however, we shall see that in fact they can be expressed as rational functions in $q, Q_1, \dots, Q_r$ which depend only on λ . The expression we obtain is a generalization of the hook length formula of Frame,

Robinson and Thrall [14] for the dimensions of the irreducible representations of the symmetric groups. This is not unexpected because $\mathcal{H} \cong RW_{r,n}$ when $R = \mathbb{C}$, $q = 1$ and $Q_s = \zeta^s$, so the orthogonality relations for the characters of $W_{r,n}$ imply that $\chi^\lambda(1) = |W_{r,n}|/s_\lambda(1; \zeta^s)$. As the referee remarked, it is worth noting that in general $P_{\mathcal{H}}(1; \zeta^s)$ is only a scalar multiple of $|W_{r,n}| = r^n n!$ since $\prod_{1 \leq i < j \leq r} (\zeta^i - \zeta^j) = (-1)^{\frac{1}{2}(r-1)} e^{\frac{r}{2}}$ by [19, 2.22].

We will compute the Schur elements of $\mathcal{H}$ by explicitly constructing a set of primitive idempotents in $\mathcal{H}$ and then applying the following Lemma (which is really a well-known fact about symmetric algebras).

2.6. Lemma. *Assume that R is a field and that $\mathcal{H}$ is semisimple. Let λ be a multipartition of n and suppose that e_λ is a primitive idempotent in $\mathcal{H}$ such that $S^\lambda \cong e_\lambda \mathcal{H}$. Then $s_\lambda(\mathbf{q}) = \frac{1}{\tau(e_\lambda)}$.*

Proof. Suppose first that R is a field of characteristic zero. Let E_λ be the primitive central idempotent corresponding to the irreducible module S^λ . By definition $\tau = \sum_{\mu} \frac{1}{s_\mu(\mathbf{q})} \chi^\mu$, where μ runs over the multipartitions of n ; therefore,

$$\tau(E_\lambda) = \sum_{\mu \vdash n} \frac{1}{s_\mu(\mathbf{q})} \chi^\mu(E_\lambda) = \frac{1}{s_\lambda(\mathbf{q})} \chi^\lambda(E_\lambda) = \frac{1}{s_\lambda(\mathbf{q})} \chi^\lambda(1).$$

Now $E_\lambda = e_1 + \dots + e_N$, where $e_1, \dots, e_N$ are primitive idempotents with $e_i \mathcal{H} \cong S^\lambda$, for all i , and $N = \dim S^\lambda = \chi^\lambda(1)$. The idempotents $e_1, \dots, e_N$ belong to the same Wedderburn component of $\mathcal{H}$, so there exist invertible elements $u_i \in \mathcal{H}$ such that $e_i = u_i e_1 u_i^{-1}$ for all i . Consequently, $\tau(e_a) = \tau(u_i e_1 u_i^{-1}) = \tau(e_1)$ since τ is a trace form; hence, $\tau(E_\lambda) = \chi^\lambda(1) \tau(e_1)$. Without loss, $e_1 = e_\lambda$ so the Lemma follows.

The case where R is a field of positive characteristic now follows by a specialization argument (using, for example, Theorem 3.14) which we leave to the reader. $\square$

2.7. Example. Fix t with $1 \leq t \leq r$ and let $\eta_t = (\eta_t^{(1)}, \dots, \eta_t^{(r)})$ be the multipartition of n with $\eta_t^{(s)} = (n)$ if $s = t$ and $\eta_t^{(s)} = (0)$ otherwise. We will compute the Schur elements $s_{\eta_t}(\mathbf{q})$. Let $x_{\eta_t} = \sum_{w \in \mathfrak{S}_n} T_w$ and $u_{\eta_t} = \prod_{s \neq t} \prod_{k=1}^n (L_k - Q_s)$ — in the product, $1 \leq s \leq r$ — and set $m_{\eta_t} = u_{\eta_t} x_{\eta_t} = x_{\eta_t} u_{\eta_t}$ (cf. (3.1)). It follows from (2.1) that u_{η_t} is central in $\mathcal{H}$. Further, the relations imply that $T_0 u_{\eta_t} = Q_t u_{\eta_t}$ and $T_w x_{\eta_t} = q^{\ell(w)} x_{\eta_t}$ for $w \in \mathfrak{S}_n$; it follows that $L_k m_{\eta_t} = q^{k-1} Q_t m_{\eta_t}$ for $k = 1, \dots, n$. Therefore, the module $m_{\eta_t} \mathcal{H}$ is one dimensional and, in particular, irreducible; in fact, $S^{\eta_t} \cong m_{\eta_t} \mathcal{H} = R m_{\eta_t}$ (for example, use (3.2)). Moreover, by what we have said

$$m_{\eta_t}^2 = [n]_q! \prod_{s \neq t} \prod_{k=1}^n (q^{k-1} Q_t - Q_s) \cdot m_{\eta_t},$$

where $[n]_q! = \prod_{k=1}^n (1 + q + \dots + q^{k-1})$; so m_{η_t} is a scalar multiple of the primitive idempotent which generates S^{η_t} . Hence, by the Lemma,

$$\begin{aligned} s_{\eta_t}(\mathbf{q}) &= (\tau(m_{\eta_t}))^{-1} [n]_q! \prod_{s \neq t} \prod_{k=0}^{n-1} (q^k Q_t - Q_s) \\ &= (-1)^{n(r-1)} [n]_q! \prod_{s \neq t} Q_s^{-n} \cdot \prod_{s \neq t} \prod_{k=0}^{n-1} (q^k Q_t - Q_s). \end{aligned}$$

Similar arguments give the Schur elements for the multipartition which is conjugate to η_t ; alternatively, they are given by Corollary 4.5 and the calculation above.

There is an action of $\mathfrak{S}_r$ on the set of multipartitions of n (by permuting components) and also on the rational functions in $Q_1, \dots, Q_r$ (by permuting parameters). When $\mathcal{H}$ is semisimple the Specht modules are determined up to isomorphism by the action of $L_1, \dots, L_n$; as the relation $\prod_{s=1}^r (T_0 - Q_s) = 0$ is invariant under the $\mathfrak{S}_r$ -action it follows that $s_{v \cdot \lambda}(\mathbf{q}) = v \cdot s_\lambda(\mathbf{q})$ for all multipartitions λ and all $v \in \mathfrak{S}_r$; this is also clear from Theorem 3.14(i). In the case where $\lambda = \eta_t$ this symmetry is evident in the formulae above.

3. AN ORTHOGONAL BASIS FOR $\mathcal{H}$

If R is a field and $P_{\mathcal{H}}(\mathbf{q})$ is non-zero then $\mathcal{H}$ is a split semisimple algebra; hence, $\mathcal{H}$ has a basis which corresponds to the matrix units in its Wedderburn decomposition. In this section we explicitly construct a Wedderburn basis for $\mathcal{H}$. We begin by recalling the standard basis of $\mathcal{H}$ from [12].

A multipartition of n (with r components) is an r -tuple $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ of partitions such that $|\lambda^{(1)}| + \dots + |\lambda^{(r)}| = n$. Let Λ^+ be the set of multipartitions of n ; then Λ^+ becomes a poset under dominance where $\lambda \supseteq \mu$ if for all $1 \leq s \leq r$ and all $i \geq 1$ we have

$$\sum_{t=1}^{s-1} |\lambda^{(t)}| + \sum_{j=1}^i \lambda_j^{(s)} \geq \sum_{t=1}^{s-1} |\mu^{(t)}| + \sum_{j=1}^i \mu_j^{(s)}.$$

We also write $\lambda \triangleright \mu$ if $\lambda \supseteq \mu$ and $\lambda \neq \mu$.

The diagram of a multipartition λ is the set

$$[\lambda] = \{ (i, j, c) \mid 1 \leq j \leq \lambda_i^{(c)} \text{ and } 1 \leq c \leq r \}.$$

We will think of $[\lambda]$ as being the r -tuple of diagrams of the partitions $\lambda^{(c)}$, for $1 \leq c \leq r$. A λ -tableau is a bijection $\mathbf{t}: [\lambda] \rightarrow \{1, 2, \dots, n\}$. If $\mathbf{t}$ is a λ -tableau write $\text{Shape}(\mathbf{t}) = \lambda$. As with diagrams, we will think of a tableau $\mathbf{t}$ as an r -tuple of tableaux $\mathbf{t} = (\mathbf{t}^{(1)}, \dots, \mathbf{t}^{(r)})$, where $\mathbf{t}^{(c)}$ is a $\lambda^{(c)}$ -tableau. The tableaux $\mathbf{t}^{(c)}$ are called the components of $\mathbf{t}$. A tableau is standard if in each component the entries increase along the rows and down the columns; let $\text{Std}(\lambda)$ be the set of standard λ -tableaux.

We identify a tableau $\mathbf{t}$ with an r -tuple of labelled diagrams; for example

$$\left(\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 5 & 6 \\ \hline 7 & \\ \hline \end{array}, \begin{array}{|c|} \hline 8 \\ \hline 9 \\ \hline \end{array} \right) \quad \text{and} \quad \left(\begin{array}{|c|c|c|} \hline 1 & 3 & 6 \\ \hline 2 & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 4 & 9 \\ \hline 7 & \\ \hline \end{array}, \begin{array}{|c|} \hline 5 \\ \hline 8 \\ \hline \end{array} \right)$$

are two $((3, 1), (2, 1), (1^2))$ -tableaux. Both of these tableaux are standard.

Given a multipartition λ let $\mathbf{t}^\lambda$ be the λ -tableau with the numbers $1, 2, \dots, n$ entered in order first along the rows of $\mathbf{t}^{\lambda^{(1)}}$ and then the rows of $\mathbf{t}^{\lambda^{(2)}}$ and so on. For example, the first of the $((3, 1), (2, 1), (1^2))$ -tableaux above is $\mathbf{t}^{((3, 1), (2, 1), (1^2))}$.

The symmetric group $\mathfrak{S}_n$ acts from the right on the set of λ -tableaux; let $\mathfrak{S}_\lambda = \mathfrak{S}_{\lambda^{(1)}} \times \dots \times \mathfrak{S}_{\lambda^{(r)}}$ be the row stabilizer of $\mathbf{t}^\lambda$. For any λ -tableau $\mathbf{t}$ let $d(\mathbf{t})$ be the unique element of $\mathfrak{S}_n$ such that $\mathbf{t} = \mathbf{t}^\lambda d(\mathbf{t})$.

Let $*$ be the R -linear antiautomorphism of $\mathcal{H}$ determined by $T_i^* = T_i$ for all i with $0 \leq i \leq n-1$. Then $T_w^* = T_{w^{-1}}$ and $L_k^* = L_k$ for all $w \in \mathfrak{S}_n$ and for $k = 1, 2, \dots, n$.

We can now define the cellular basis of $\mathcal{H}$ constructed in [12]. Fix a multipartition λ and let $\mathbf{a} = (\mathbf{a}_1, \dots, \mathbf{a}_r)$, where $\mathbf{a}_s = |\lambda^{(1)}| + \dots + |\lambda^{(s-1)}|$ for $1 \leq s \leq r$. Define $m_\lambda = x_\lambda u_\lambda^+$, where

$$x_\lambda = \sum_{w \in \mathfrak{S}_\lambda} T_w \quad \text{and} \quad u_\lambda^+ = \prod_{s=2}^r \prod_{k=1}^{\mathbf{a}_s} (L_k - Q_s).$$

It follows from (2.1) that x_λ and u_λ^+ commute. Finally, given standard λ -tableaux $\mathfrak{s}$ and $\mathfrak{t}$ let $m_{\mathfrak{s}\mathfrak{t}} = T_{d(\mathfrak{s})}^* m_\lambda T_{d(\mathfrak{t})}$.

Whenever we write $m_{\mathfrak{s}\mathfrak{t}}$ in what follows $\mathfrak{s}$ and $\mathfrak{t}$ will be standard tableaux of the same shape (and similarly, for $f_{\mathfrak{s}\mathfrak{t}}$ etc.).

3.1 (Dipper–James–Mathas [12, Theorem 3.26]) *The Ariki–Koike algebra $\mathcal{H}$ is free as an R -module with cellular basis $\{m_{\mathfrak{s}\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \text{Std}(\lambda) \text{ for some } \lambda \in \Lambda^+\}$.*

One consequence of this result is that the R -module $\check{\mathcal{H}}^\lambda$ which has as basis the set of $m_{\mathfrak{u}\mathfrak{v}}$ where $\text{Shape}(\mathfrak{u}) = \text{Shape}(\mathfrak{v}) \triangleright \lambda$ is a two-sided ideal of $\mathcal{H}$. It follows from (3.1) that $m_\mu \mathcal{H} \subset \check{\mathcal{H}}^\lambda$ whenever $\mu \triangleright \lambda$. The Specht module S^λ is the right $\mathcal{H}$ -module $m_\lambda \mathcal{H} / (m_\lambda \mathcal{H} \cap \check{\mathcal{H}}^\lambda)$, a submodule of $\mathcal{H} / \check{\mathcal{H}}^\lambda$. Thus, S^λ is free as an R -module with basis $\{m_{\mathfrak{t}} \mid \mathfrak{t} \in \text{Std}(\lambda)\}$, where $m_{\mathfrak{t}} = m_{\mathfrak{t}\lambda} + \check{\mathcal{H}}^\lambda$. Further, there is a natural associative bilinear form $\langle \cdot, \cdot \rangle$ on S^λ which is determined by

$$\langle m_{\mathfrak{s}}, m_{\mathfrak{t}} \rangle m_\lambda \equiv m_{\mathfrak{t}\lambda} m_{\mathfrak{s}\lambda} \pmod{\check{\mathcal{H}}^\lambda}.$$

Before we can begin we need some further notation and a result from [17]. If $\mathfrak{t}$ is any tableau and $k \geq 0$ is an integer let $\mathfrak{t} \downarrow k$ be the subtableau of $\mathfrak{t}$ which contains the integers $1, 2, \dots, k$. Observe that $\mathfrak{t}$ is standard if and only if $\text{Shape}(\mathfrak{t} \downarrow k)$ is a multipartition for all k . We extend the dominance order to the set of standard tableaux by defining $\mathfrak{s} \supseteq \mathfrak{t}$ if $\text{Shape}(\mathfrak{s} \downarrow k) \supseteq \text{Shape}(\mathfrak{t} \downarrow k)$ for $k = 1, 2, \dots, n$; again, we write $\mathfrak{s} \triangleright \mathfrak{t}$ if $\mathfrak{s} \supseteq \mathfrak{t}$ and $\mathfrak{s} \neq \mathfrak{t}$.

Write $\text{res}_i(k) = q^{j-i} Q_c$ if k appears in row i and column j of $\mathfrak{t}^{(c)}$; then $\text{res}_{\mathfrak{t}}(k)$ is the residue of k in $\mathfrak{t}$. We can now state the result we need from [17].

3.2 [17, Prop. 3.7] *Let $\mathfrak{s}$ and $\mathfrak{t}$ be standard λ -tableaux and suppose that k is an integer with $1 \leq k \leq n$. Then there exist $a_{\mathfrak{v}} \in R$ such that*

$$m_{\mathfrak{s}\mathfrak{t}} L_k = \text{res}_{\mathfrak{t}}(k) m_{\mathfrak{s}\mathfrak{t}} + \sum_{\substack{\mathfrak{v} \in \text{Std}(\lambda) \\ \mathfrak{v} \triangleright \mathfrak{t}}} a_{\mathfrak{v}} m_{\mathfrak{s}\mathfrak{v}} \pmod{\check{\mathcal{H}}^\lambda}.$$

From our current point of view the importance of this result derives from the observation that $m_{\mathfrak{s}\mathfrak{t}}(L_k - \text{res}_{\mathfrak{u}}(k)) / (\text{res}_{\mathfrak{t}}(k) - \text{res}_{\mathfrak{u}}(k)) = m_{\mathfrak{s}\mathfrak{t}}$ plus a linear combination of more dominant terms, providing that $\text{res}_{\mathfrak{u}}(k) \neq \text{res}_{\mathfrak{t}}(k)$; this motivates the next definition.

Let $\mathcal{R}(k) = \{q^d Q_s \mid 1 \leq s \leq r, |d| < k \text{ and } d \neq 0 \text{ if } r = 1 \text{ and } k = 2, 3\}$; be the set of possible residues $\text{res}_{\mathfrak{t}}(k)$ as $\mathfrak{t}$ runs over the standard tableaux.

3.3. Definition (cf. [17, Defn. 3.11]). *Suppose that $\mathfrak{s}$ and $\mathfrak{t}$ are standard λ -tableaux.*

- (i) *Let $F_{\mathfrak{t}} = \prod_{k=1}^n \prod_{\substack{c \in \mathcal{R}(k) \\ c \neq \text{res}_{\mathfrak{t}}(k)}} \frac{L_k - c}{\text{res}_{\mathfrak{t}}(k) - c}$.*
- (ii) *Let $f_{\mathfrak{s}\mathfrak{t}} = F_{\mathfrak{s}} m_{\mathfrak{s}\mathfrak{t}} F_{\mathfrak{t}}$.*

There are some remarks worth making about the definition of $F_{\mathfrak{t}}$. First, we do not need to specify an order for the product in the definition of $F_{\mathfrak{t}}$ since the L_k

generate a commutative subalgebra of $\mathcal{H}$. Secondly, the definition of F_t is very conservative in the sense that many of the factors of F_t can be omitted without changing the element f_{st} . Finally, historically this construction has been used to produce an orthogonal basis for the Specht modules; we are going to modify this procedure to give an orthogonal basis for the whole of $\mathcal{H}$.

Let $\mathfrak{s}$ and $\mathfrak{t}$ be two standard tableaux, not necessarily of the same shape. The proof of next result rests upon the easy fact [17, Lemma 3.12] that, because $P_{\mathcal{H}}(\mathbf{q})$ is invertible, $\mathfrak{s} = \mathfrak{t}$ if and only if $\text{res}_{\mathfrak{s}}(k) = \text{res}_{\mathfrak{t}}(k)$ for $k = 1, \dots, n$.

3.4. Proposition. *Suppose that $P_{\mathcal{H}}$ is invertible in R and that $\mathfrak{s}$ and $\mathfrak{t}$ are standard λ -tableau and that k is an integer with $1 \leq k \leq n$. Then*

- (i) $f_{st} = m_{st} + \sum_{(u,v) \triangleright (s,t)} a_{uv} m_{uv}$ for some $a_{uv} \in R$;
- (ii) $f_{st} L_k = \text{res}_{\mathfrak{t}}(k) f_{st}$;
- (iii) $f_{st} F_u = \delta_{tu} f_{st}$; and,
- (iv) $F_u f_{st} = \delta_{su} f_{st}$.

Proof. Given (3.2), this is a variation on a well known argument; see, for example, [22, Prop. 3.35]. The proof in [22] can be copied out verbatim except that N should be replaced by $\sum_{k=1}^n |\mathcal{R}(k)|$. $\square$

In particular, part (i) together with (3.1) shows that

$$\{ f_{st} \mid \mathfrak{s}, \mathfrak{t} \in \text{Std}(\lambda) \text{ for some } \lambda \vdash n \}$$

is a basis of $\mathcal{H}$; shortly we will see that it is an orthogonal basis of $\mathcal{H}$ with respect to the trace form τ . As a first step we describe the action of $\mathcal{H}(\mathfrak{S}_n)$ on this basis; note that the action of the L_k (and, in particular, $T_0 = L_1$), on this basis is given by Proposition 3.4(ii).

3.5. Proposition. *Suppose that $\mathfrak{t} = \mathfrak{s}(i, i+1)$ where $\mathfrak{s}$ and $\mathfrak{u}$ are standard λ -tableaux and i is an integer with $1 \leq i < n$. If $\mathfrak{t}$ is standard then*

$$f_{us} T_i = \begin{cases} \frac{(q-1) \text{res}_{\mathfrak{t}}(i)}{\text{res}_{\mathfrak{t}}(i) - \text{res}_{\mathfrak{s}}(i)} f_{us} + f_{ut}, & \text{if } \mathfrak{s} \triangleright \mathfrak{t}, \\ \frac{(q-1) \text{res}_{\mathfrak{t}}(i)}{\text{res}_{\mathfrak{t}}(i) - \text{res}_{\mathfrak{s}}(i)} f_{us} + \frac{(q \text{res}_{\mathfrak{s}}(i) - \text{res}_{\mathfrak{t}}(i))(\text{res}_{\mathfrak{s}}(i) - q \text{res}_{\mathfrak{t}}(i))}{(\text{res}_{\mathfrak{t}}(i) - \text{res}_{\mathfrak{s}}(i))^2} f_{ut}, & \text{if } \mathfrak{t} \triangleright \mathfrak{s}. \end{cases}$$

If $\mathfrak{t}$ is not standard then

$$f_{us} T_i = \begin{cases} q f_{us}, & \text{if } i \text{ and } i+1 \text{ are in the same row of } \mathfrak{s}, \\ -f_{us}, & \text{if } i \text{ and } i+1 \text{ are in the same column of } \mathfrak{s}. \end{cases}$$

Proof. By the above remarks, $\{f_{us}\}$ is a basis of $\mathcal{H}$, so $f_{us} T_i = \sum_{\mathfrak{a}, \mathfrak{b}} r_{\mathfrak{a}\mathfrak{b}} f_{\mathfrak{a}\mathfrak{b}}$ for some $r_{\mathfrak{a}\mathfrak{b}} \in R$. By Proposition 3.4(iv), $F_u f_{\mathfrak{a}\mathfrak{b}} = \delta_{\mathfrak{a}u} f_{\mathfrak{u}\mathfrak{b}}$. Therefore, multiplying the equation for $f_{us} T_i$ on the left by F_u shows that $r_{\mathfrak{a}\mathfrak{b}} = 0$ whenever $\mathfrak{a} \neq u$; in particular, $r_{\mathfrak{a}\mathfrak{b}} = 0$ if $\text{Shape}(\mathfrak{b}) \neq \lambda$. Hence, $f_{us} T_i = \sum_{\mathfrak{b}} a_{\mathfrak{b}} f_{\mathfrak{u}\mathfrak{b}}$, for some $a_{\mathfrak{b}} \in R$, where $\mathfrak{b}$ runs over the set of standard λ -tableaux. The argument of [22, Theorem 3.36] can now be repeated, essentially word for word, to complete the proof. $\square$

For each standard λ -tableau $\mathfrak{s}$ let $f_{\mathfrak{s}} = f_{t^{\lambda}_{\mathfrak{s}}} + \check{\mathcal{H}}^{\lambda}$. Then $\{f_{\mathfrak{s}} \mid \mathfrak{s} \in \text{Std}(\lambda)\}$ is a basis of the Specht module S^{λ} by Proposition 3.4(i). Note that $f_{t^{\lambda}} = m_{t^{\lambda}}$; further, by Proposition 3.5, if $\mathfrak{s}$ and $\mathfrak{t}$ are standard λ -tableaux with $\mathfrak{s} = \mathfrak{t}(i, i+1) \triangleright \mathfrak{t}$ then $f_{\mathfrak{t}} = f_{\mathfrak{s}}(T_i - \alpha)$, where $\alpha = (q-1) \text{res}_{\mathfrak{t}}(i) / (\text{res}_{\mathfrak{t}}(i) - \text{res}_{\mathfrak{s}}(i))$; hence, by [17,

Cor. 3.14], $\{f_{\mathfrak{s}} \mid \mathfrak{s} \in \text{Std}(\lambda)\}$ is the orthogonal basis of S^λ which was constructed in [17, Theorem 3.13(vi)]. (In the notation of [17], $f_{\mathfrak{s}} = \psi_{\mathfrak{s}}(m_\lambda)$.)

The inner products $\langle f_{\mathfrak{s}}, f_{\mathfrak{t}} \rangle$, for $\mathfrak{s}, \mathfrak{t} \in \text{Std}(\lambda)$ were computed explicitly (as rational functions) in [17]; to describe this we need some more notation. Recall that a node $y \notin [\lambda]$ is an **addable node** for λ if $[\lambda] \cup \{y\}$ is the diagram of a multipartition; similarly, $y \in [\lambda]$ is a **removable node** for λ if $[\lambda] \setminus \{y\}$ is the diagram of a multipartition. Given two nodes $x = (i, j, k)$ and $y = (a, b, c)$, write $y < x$ if either $c < k$, or $c = k$ and $b > j$.

Let $\mathfrak{s}$ be a λ -tableau. Then for each integer i there is a unique node $x \in [\lambda]$ such that $\mathfrak{s}(x) = i$, $1 \leq i \leq n$. Let $\mathcal{A}_{\mathfrak{s}}(i)$ be the set of addable nodes for $\text{Shape}(\mathfrak{s} \downarrow i)$ which are strictly less than x (with respect to $<$); similarly, let $\mathcal{R}_{\mathfrak{s}}(i)$ be the set of removable nodes strictly less than x for the multipartition $\text{Shape}(\mathfrak{s} \downarrow i - 1)$. Essentially as in [17, Defn. 3.15] let

$$(3.6) \quad \gamma_{\mathfrak{s}} = q^{\ell(d(\mathfrak{s})) + \alpha(\lambda)} \prod_{i=1}^n \frac{\prod_{y \in \mathcal{A}_{\mathfrak{s}}(i)} (\text{res}_{\mathfrak{s}}(i) - \text{res}(y))}{\prod_{y \in \mathcal{R}_{\mathfrak{s}}(i)} (\text{res}_{\mathfrak{s}}(i) - \text{res}(y))}$$

where $\alpha(\lambda) = \frac{1}{2} \sum_{s=1}^r \sum_{i \geq 1} (\lambda_i^{(s)} - 1) \lambda_i^{(s)}$.

If k is an integer let $[k]_q = \frac{q^k - 1}{q - 1}$ if $q \neq 1$; more generally, set $[k]_q = 1 + q + \dots + q^{k-1}$ if $k \geq 0$ and $[k]_q = -q^k [-k]_q$ if $k < 0$. When $k \geq 0$ we also set $[k]_q! = [1]_q [2]_q \dots [k]_q$. Finally, if λ is a multipartition let $[\lambda]_q! = \prod_{s=1}^r \prod_{i \geq 1} [\lambda_i^{(s)}]_q!$.

3.7 (James–Mathas [17, (3.17)–(3.19)]) Suppose that $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R and let $\mathfrak{t} \in \text{Std}(\lambda)$. Then $\gamma_{\mathfrak{t}}$ is uniquely determined by the two conditions

- (i) $\gamma_{\mathfrak{t}\lambda} = [\lambda]_q! \prod_{1 \leq s < t \leq r} \prod_{(i,j) \in [\lambda^{(s)}]} (q^{j-i} Q_s - Q_t)$; and,
- (ii) if $\mathfrak{s} = \mathfrak{t}(i, i+1) \triangleright \mathfrak{t}$ then $\gamma_{\mathfrak{t}} = \frac{(q \text{res}_{\mathfrak{s}}(i) - \text{res}_{\mathfrak{t}}(i))(\text{res}_{\mathfrak{s}}(i) - q \text{res}_{\mathfrak{t}}(i))}{(\text{res}_{\mathfrak{s}}(i) - \text{res}_{\mathfrak{t}}(i))^2} \gamma_{\mathfrak{s}}$.

Furthermore, $\langle f_{\mathfrak{s}}, f_{\mathfrak{t}} \rangle = \delta_{\mathfrak{s}\mathfrak{t}} \gamma_{\mathfrak{t}}$ for all $\mathfrak{s}, \mathfrak{t} \in \text{Std}(\lambda)$.

- 3.8. Remarks.** (i) In [17] the rational functions $\gamma_{\mathfrak{t}}$ were important only up to a power of q (which is a unit in R). In this paper we need to know the inner products $\langle f_{\mathfrak{s}}, f_{\mathfrak{t}} \rangle$ exactly; for this reason our definition of $\gamma_{\mathfrak{s}}$ differs from that in [17] by the factor $q^{\ell(d(\mathfrak{s})) + \alpha(\lambda)}$. The argument in [17] computes $\langle f_{\mathfrak{s}}, f_{\mathfrak{t}} \rangle$ using only (3.7)(i) and (3.7)(ii); the inner product $\langle f_{\mathfrak{t}\lambda}, f_{\mathfrak{t}\lambda} \rangle$ is easily seen to be given by the formula on the right hand side of (3.7)(i).
- (ii) A factor of q was omitted from the formula for $\gamma_{\mathfrak{t}}$ in [17, Lemma 3.17]; this is corrected in (3.7)(ii) — note that $\ell(d(\mathfrak{t})) = \ell(d(\mathfrak{s})) + 1$.
- (iii) The definition of $\gamma_{\mathfrak{s}}$ above is simpler than that given in [17] because $\mathfrak{s}$ is a standard tableau (rather than the more general semistandard tableau which were considered in [17]).

Let $(,)$ be the inner product on $\mathcal{H}$ given by $(h_1, h_2) = \tau(h_1 h_2^*)$, for $h_1, h_2 \in \mathcal{H}$. Then $(,)$ is a symmetric associative bilinear form on $\mathcal{H}$.

3.9. Theorem. Suppose that $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R . Then

$$\{f_{\mathfrak{s}\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \text{Std}(\lambda) \text{ for some } \lambda \in \Lambda^+\}$$

is a orthogonal basis of $\mathcal{H}$ with respect to the trace form τ . In addition, if $\mathfrak{s}, \mathfrak{t}, \mathfrak{u}$ and $\mathfrak{v}$ are standard tableaux then $f_{\mathfrak{s}\mathfrak{t}} f_{\mathfrak{u}\mathfrak{v}} = \delta_{\mathfrak{u}\mathfrak{t}} \gamma_{\mathfrak{t}} f_{\mathfrak{s}\mathfrak{v}}$.

Proof. By (3.1) the set $\{m_{st}\}$ is a basis of $\mathcal{H}$; therefore, $\{f_{st}\}$ is also a basis of $\mathcal{H}$ by Proposition 3.4(i). Next we prove that $f_{st}f_{uv} = \delta_{ut}\gamma_t f_{sv}$. First, if $u \neq t$ then $f_{st}f_{uv} = f_{st}F_u m_{uv} F_v = 0$ by Proposition 3.4(iii). Now consider $f_{st}f_{tv}$; since $\{f_{ab}\}$ is a basis we can write $f_{st}f_{tv} = \sum_{a,b} r_{ab}^t f_{ab}$ for some $r_{ab}^t \in R$. Applying Proposition 3.4(iii) and its left handed analogue shows that

$$f_{st}f_{tv} = F_s f_{st}f_{tv} F_v = \sum_{a,b} r_{ab}^t F_s f_{ab} F_v = r_{sv}^t f_{sv}.$$

Now, for any $s, v \in \text{Std}(\lambda)$ the definition of the inner product on the Specht module gives

$$\langle f_t, f_t \rangle f_{sv} \equiv f_{st}f_{tv} \pmod{\tilde{\mathcal{H}}^\lambda}.$$

Therefore, $r_{sv}^t = \langle f_t, f_t \rangle = \gamma_t$ by (3.7); so r_{sv}^t depends only on t and $f_{st}f_{tv} = \gamma_t f_{sv}$ as claimed.

Finally, it remains to show that the basis $\{f_{st}\}$ is orthogonal with respect to the bilinear form $(\ , \)$. First, $(f_{st}, f_{uv}) = \tau(f_{st}f_{uv}^*) = \tau(f_{st}f_{vu}) = \delta_{tv}\gamma_t\tau(f_{su})$; in particular, $(f_{st}, f_{uv}) = 0$ if $t \neq v$. On the other hand, τ is a trace form so $(f_{st}, f_{uv}) = \tau(f_{st}f_{vu}) = \tau(f_{vu}f_{st}) = \delta_{su}\gamma_s\tau(f_{vt})$. Therefore, $(f_{st}, f_{uv}) = \delta_{su}\delta_{tv}\gamma_s\tau(f_{tt})$. Consequently, $\{f_{st}\}$ is an orthogonal basis of $\mathcal{H}$ (and $\tau(f_{tt})$ is non-zero for all t). $\square$

The basis $\{f_{st}\}$ is cellular (with respect to the involution $*$); but this is not surprising as it was constructed from a cellular basis.

3.10. Remark. In [21] it was shown that if $\mathcal{H}$ is defined over a ring R in which the parameters $q, Q_1, \dots, Q_r$ are invertible then $\mathcal{H}$ is a symmetric algebra with respect to the trace form τ ; however, this was proved indirectly without constructing a pair of dual bases. The Theorem gives a self-dual basis of the semisimple Ariki-Koike algebras; no such basis is known in general.

As a first consequence, Theorem 3.9 identifies a submodule of $\mathcal{H}$ which is isomorphic to the Specht module S^λ .

3.11. Corollary. *Suppose that $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R and let s and t be standard λ -tableaux. Then $S^\lambda \cong f_{st}\mathcal{H} = \sum_{v \in \text{Std}(\lambda)} R f_{sv}$.*

Proof. By the Theorem, $f_{st}\mathcal{H}$ has as basis the set $\{f_{sv} \mid v \in \text{Std}(\lambda)\}$. The isomorphism is given by the linear map $S^\lambda \rightarrow f_{st}\mathcal{H}$ determined by $f_v \mapsto f_{sv}$ for all tableaux $v \in \text{Std}(\lambda)$. $\square$

Set $\tilde{f}_{st} = \gamma_t^{-1} f_{st}$. Then $\tilde{f}_{st}\tilde{f}_{uv} = \delta_{tu}\tilde{f}_{sv}$ and $\{\tilde{f}_{st}\}$ is a basis of $\mathcal{H}$. Hence, we have the following.

3.12. Corollary. *Suppose that $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R . Then*

$$\{\tilde{f}_{st} \mid s, t \in \text{Std}(\lambda) \text{ for some } \lambda \in \Lambda^+\}$$

is a basis of matrix units in $\mathcal{H}$.

The last result yields an explicit isomorphism from $\mathcal{H}$ to the group ring of $W_{r,n}$ when $P_{\mathcal{H}}(\mathbf{q})$ is invertible. Assume that R contains a primitive r th root of unity ζ ; then, $\mathcal{H} \cong RW_{r,n}$ when $q = 1$ and $Q_s = \zeta^s$ for $s = 1, 2, \dots, r$. Write $\tilde{f}_{st}^1$ for the element of $RW_{r,n}$ corresponding to $\tilde{f}_{st} \in \mathcal{H}$ under the canonical isomorphism $\mathcal{H} \rightarrow RW_{r,n}$.

3.13. Corollary. *Assume that R contains a primitive r th root of unity and that $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R . Then $\mathcal{H} \cong RW_{r,n}$ via the R -algebra homomorphism determined by $\tilde{f}_{\mathbf{st}} \mapsto \tilde{f}_{\mathbf{st}}^1$.*

By parts (i) and (iii) of Theorem 3.14 below, $T_i = \sum_{\mathbf{t}} \tilde{f}_{\mathbf{t}} T_i$, for $0 \leq i < n$; so, in principle, we can determine the image of the generators of $\mathcal{H}$ under this isomorphism.

Lusztig [18] has shown that there exists a homomorphism Φ from the Hecke algebra $\mathcal{H}(W)$ of any finite Weyl group W to the group ring RW and he shows that Φ induces an isomorphism when $\mathcal{H}(W)$ is semisimple. Our map is not an analogue of Lusztig's isomorphism; rather it is an explicit realization of the Tits deformation theorem in this setting. It would be good to find a generalization of Lusztig's isomorphism theorem for the Ariki–Koike algebras.

We next construct the primitive (central) idempotents in $\mathcal{H}$.

3.14. Theorem. *Suppose that R is a field and that $P_{\mathcal{H}}(\mathbf{q}) \neq 0$.*

- (i) *Let $\mathbf{t}$ be a standard λ -tableau. Then $F_{\mathbf{t}} = \frac{1}{\gamma_{\mathbf{t}}} f_{\mathbf{t}\mathbf{t}}$ and $F_{\mathbf{t}}$ is a primitive idempotent with $S^{\lambda} \cong F_{\mathbf{t}}\mathcal{H}$.*
- (ii) *For any multipartition λ let $F_{\lambda} = \sum_{\mathbf{t} \in \text{Std}(\lambda)} F_{\mathbf{t}}$. Then F_{λ} is a primitive central idempotent.*
- (iii) *$\{F_{\lambda} \mid \lambda \vdash n\}$ is a complete set of primitive central idempotents; in particular,*

$$1 = \sum_{\lambda \vdash n} F_{\lambda} = \sum_{\mathbf{t} \text{ standard}} F_{\mathbf{t}}.$$

Proof. We may write $F_{\mathbf{t}} = \sum_{\mathbf{u}, \mathbf{v}} a_{\mathbf{u}\mathbf{v}} f_{\mathbf{u}\mathbf{v}}$ for some $a_{\mathbf{u}\mathbf{v}} \in R$ by Theorem 3.9. By Proposition 3.4(iii) and Theorem 3.9, $f_{\mathbf{st}} F_{\mathbf{t}} = \sum_{\mathbf{u}, \mathbf{v}} a_{\mathbf{u}\mathbf{v}} f_{\mathbf{st}} f_{\mathbf{u}\mathbf{v}} = \sum_{\mathbf{v}} \gamma_{\mathbf{t}} a_{\mathbf{t}\mathbf{v}} f_{\mathbf{sv}}$; equating coefficients on both sides shows that $a_{\mathbf{t}\mathbf{v}} = 0$ if $\mathbf{v} \neq \mathbf{t}$ and that $1 = a_{\mathbf{t}\mathbf{t}} \gamma_{\mathbf{t}}$. Since $F_{\mathbf{t}}^* = F_{\mathbf{t}}$ we also have that $a_{\mathbf{v}\mathbf{t}} = 0$ if $\mathbf{v} \neq \mathbf{t}$. Hence, $F_{\mathbf{t}} = \frac{1}{\gamma_{\mathbf{t}}} f_{\mathbf{t}\mathbf{t}}$ as claimed. By Theorem 3.9, $\frac{1}{\gamma_{\mathbf{t}}} f_{\mathbf{t}\mathbf{t}}$ is idempotent; further, it is primitive because S^{λ} is irreducible and $S^{\lambda} \cong \frac{1}{\gamma_{\mathbf{t}}} f_{\mathbf{t}\mathbf{t}}\mathcal{H}$ by Corollary 3.11. Hence, (i) is proved.

Parts (ii) and (iii) now follow because $\mathcal{H} = \bigoplus_{\lambda \vdash n} \bigoplus_{\mathbf{t} \in \text{Std}(\lambda)} F_{\mathbf{t}}\mathcal{H}$ is a decomposition of $\mathcal{H}$ into a direct sum of simple modules with each simple module $F_{\mathbf{t}}\mathcal{H} \cong S^{\lambda}$ appearing with multiplicity equal to its dimension (the sum is direct because $\{f_{\mathbf{t}\mathbf{v}}\}$ is a basis of $F_{\mathbf{t}}\mathcal{H}$ by Theorem 3.9 and the set of all $f_{\mathbf{u}\mathbf{v}}$ is a basis of $\mathcal{H}$). $\square$

3.15. Corollary. *Let $\mathbf{t}$ be a standard tableau and let k be an integer with $1 \leq k \leq n$. Then $F_{\mathbf{t}}L_k = \text{res}_{\mathbf{t}}(k)F_{\mathbf{t}}$ and $f_{\mathbf{st}}L_k = \text{res}_{\mathbf{t}}(k)f_{\mathbf{st}}$.*

Proof. Since $F_{\mathbf{t}} = \frac{1}{\gamma_{\mathbf{t}}} f_{\mathbf{t}\mathbf{t}}$ by part (i), the formula for $F_{\mathbf{t}}L_k$ follows from Proposition 3.4(ii); this also implies the second statement because $f_{\mathbf{st}}L_k = f_{\mathbf{st}}F_{\mathbf{t}}L_k$. $\square$

In particular, the Corollary describes the action of $T_0 = L_1$ on the orthogonal basis $\{f_{\mathbf{st}}\}$ of $\mathcal{H}$. Proposition 3.5 gives the action of the remaining generators $T_1, \dots, T_{n-1}$ of $\mathcal{H}$ on the basis $\{f_{\mathbf{st}}\}$.

3.16. Corollary. *Suppose that $1 \leq k \leq n$. Then $\prod_{c \in \mathcal{R}(k)} (L_k - c) = 0$ and this is the minimum polynomial for L_k acting on $\mathcal{H}$.*

Proof. By Theorem 3.14,

$$\prod_{c \in \mathcal{R}(k)} (L_k - c) = 1 \cdot \prod_{c \in \mathcal{R}(k)} (L_k - c) = \sum_{\mathbf{t} \text{ standard}} F_{\mathbf{t}} \prod_{c \in \mathcal{R}(k)} (L_k - c) = 0,$$

since $F_{\mathbf{t}}(L_k - \text{res}_{\mathbf{t}}(k)) = 0$ by Corollary 3.15. Moreover, if we remove any factor $(L_k - c)$ from the product $\prod_{c \in \mathcal{R}(k)} (L_k - c)$ then what remains is non-zero because it divides $F_{\mathbf{t}}$ for some standard tableau $\mathbf{t}$. Hence, $\prod_{c \in \mathcal{R}(k)} (L_k - c)$ is the minimum polynomial of L_k . $\square$

3.17. Corollary. *Suppose that $1 \leq k \leq n$. Then $L_k = \sum_{\mathbf{t}} \text{res}_{\mathbf{t}}(k) F_{\mathbf{t}}$, where the sum is over the set of all standard tableaux (of arbitrary shape).*

Proof. Combining part (iii) of Theorem 3.14 with Corollary 3.15 shows that

$$L_k = 1 \cdot L_k = \sum_{\mathbf{t} \text{ standard}} F_{\mathbf{t}} L_k = \sum_{\mathbf{t} \text{ standard}} \text{res}_{\mathbf{t}}(k) F_{\mathbf{t}}$$

as required. $\square$

We close this section with a description of the centre of $\mathcal{H}$ in the semisimple case; this result is due to Ariki and Koike [4, Theorem 3.20].

3.18. Theorem (Ariki–Koike). *Suppose that R is a field and that $P_{\mathcal{H}}(\mathbf{q}) \neq 0$.*

- (i) *The centre of $\mathcal{H}$ is the set of symmetric polynomials in $L_1, L_2, \dots, L_n$.*
- (ii) *Let $\mathbb{L}$ be the subalgebra of $\mathcal{H}$ generated by $L_1, L_2, \dots, L_n$. Then $\mathbb{L}$ is a maximal abelian subalgebra of $\mathcal{H}$.*

Proof. First consider (ii). By definition $\mathbb{L}$ contains each of the primitive idempotents $F_{\mathbf{t}}$, for an arbitrary standard tableau $\mathbf{t}$. On the other hand, by Corollary 3.17, $\mathbb{L}$ is contained in the subalgebra of $\mathcal{H}$ generated by the primitive idempotents $F_{\mathbf{t}}$; hence, $\mathbb{L}$ is the subalgebra of $\mathcal{H}$ generated by the idempotents $F_{\mathbf{t}}$. As the primitive idempotents generate a maximal abelian subalgebra of $\mathcal{H}$ the result follows.

Now consider (i). By (2.1) every symmetric polynomial in $L_1, \dots, L_n$ belongs to the centre of $\mathcal{H}$. Conversely, the centre of $\mathcal{H}$ has as basis the set of idempotents F_{λ} , as λ runs over the multipartitions of n . So to prove (i) it is enough to show that each F_{λ} is symmetric in $L_1, \dots, L_n$.

Let $\mathcal{R} = \cup_{k=1}^n \mathcal{R}(k)$ be the set of all possible residues for $\mathcal{H}$. By Corollary 3.15 if $\mathbf{t}$ is a λ -tableau and $1 \leq m \leq n$ then $F_{\mathbf{t}}(L_k - c) = (\text{res}_{\mathbf{t}}(k) - c)F_{\mathbf{t}}$; therefore, multiplying $F_{\mathbf{t}}$ by the appropriate extra factors we see that

$$F_{\mathbf{t}} \stackrel{\text{def}}{=} \prod_{k=1}^n \prod_{c \in \mathcal{R}(k) \setminus \{\text{res}_{\mathbf{t}}(k)\}} \frac{L_k - c}{\text{res}_{\mathbf{t}}(k) - c} = \prod_{k=1}^n \prod_{c \in \mathcal{R} \setminus \{\text{res}_{\mathbf{t}}(k)\}} \frac{L_k - c}{\text{res}_{\mathbf{t}}(k) - c}.$$

Notice that the denominator is equal to $f_{\lambda} = \prod_{k=1}^n \prod_{c \in \mathcal{R} \setminus \{\text{res}_{\mathbf{t}}(k)\}} (\text{res}_{\mathbf{t}}(k) - c)$ and f_{λ} depends only on λ and not directly on $\mathbf{t}$.

Next, following Murphy [25] say that a λ -tableau is **regular** if its entries increase from left to right along the nodes in $[\lambda]$ of constant residue. Now, because $P_{\mathcal{H}}(\mathbf{q}) \neq 0$ two nodes in $[\lambda]$ have the same residue if and only if they lie on the same diagonal $\{(i + d, j + d, s) \in [\lambda^{(s)}] \mid d \geq 0\}$ of $[\lambda]$; thus, a tableau is regular if and only if its entries increase from left to right along each diagonal. (So, for example, every

standard tableau is regular, but not conversely.) Extending the formula above for F_t , for each regular tableau t define

$$F_t = \prod_{k=1}^n \prod_{c \in \mathcal{R} \setminus \{\text{res}_t(k)\}} \frac{L_k - c}{\text{res}_t(k) - c} = \frac{1}{f_\lambda} \prod_{k=1}^n \prod_{c \in \mathcal{R} \setminus \{\text{res}_t(k)\}} (L_k - c).$$

Observe that if $c \in \mathcal{R}$ then $L_k - c$ is a factor of F_t if and only if $c \neq \text{res}_t(k)$; therefore, F_t determines $\text{res}_t(k)$, for $k = 1, \dots, n$. As remarked above, the residues on the different diagonals of $[\lambda]$ are distinct; consequently, a regular tableau t is uniquely determined by the sequence of residues $(\text{res}_t(1), \dots, \text{res}_t(n))$ and hence by the ‘polynomial’ F_t . It follows that if we permute $L_1, \dots, L_n$ then F_t is mapped to F_s where s is the regular tableau determined by the corresponding permutation of the residue sequence of t — again, s is necessarily a λ -tableau because the shape of s is determined by the lengths of its diagonals which, in turn, are determined by the multiplicity of each residue in s (or t).

Finally, notice that a regular tableau t is *not* standard if and only if $\text{res}_t(k) \notin \mathcal{R}(k)$ for some k . Therefore, if t is not standard then $\prod_{c \in \mathcal{R}(k)} (L_k - c)$ is a factor of F_t and, consequently, $F_t = 0$ by Corollary 3.16; hence,

$$F_\lambda = \sum_{\substack{t \text{ a standard} \\ \lambda\text{-tableau}}} F_t = \sum_{\substack{t \text{ a regular} \\ \lambda\text{-tableau}}} F_t.$$

By the last paragraph the right hand side is a symmetric polynomial in $L_1, \dots, L_n$ so the theorem follows. $\square$

For the Iwahori–Hecke algebras of type A (that is, when $r = 1$), it is conjectured that the centre of $\mathcal{H}$ is always the set of symmetric polynomials in $L_1, \dots, L_n$, even when $\mathcal{H}$ is not semisimple; see [10, 23]. When $r > 1$ and $\mathcal{H}$ is not semisimple there are cases where the centre of $\mathcal{H}$ is larger than the set of symmetric polynomials in $L_1, \dots, L_n$; for an example see [3, p. 792].

4. ANOTHER CONSTRUCTION OF THE SPECHT MODULES

In this section we give another two constructions of the Specht modules. The first is via a second cellular basis of $\mathcal{H}$ which is, in a certain sense, dual to the basis described in the previous section. The second construction combines these two approaches to produce submodules of $\mathcal{H}$ which are isomorphic to the Specht modules S^λ . Some of these results can be found in the work of Du and Rui [13]. In the next section we will use these results to compute the Schur elements of $\mathcal{H}$.

Let $\mathcal{Z} = \mathbb{Z}[\hat{q}, \hat{q}^{-1}, \hat{Q}_1, \dots, \hat{Q}_r]$, where $\hat{q}, \hat{Q}_1, \dots, \hat{Q}_r$ are indeterminates over $\mathbb{Z}$, and let $\mathcal{H}_{\mathcal{Z}}$ be the Ariki–Koike algebra with parameters $\hat{q}, \hat{Q}_1, \dots, \hat{Q}_r$. Consider the ring R as a $\mathcal{Z}$ -module by letting $\hat{q}$ act on R as multiplication by q and $\hat{Q}_s$ by multiplication by Q_s , for $1 \leq s \leq r$. Then $\mathcal{H} \cong \mathcal{H}_{\mathcal{Z}} \otimes_{\mathcal{Z}} R$, since $\mathcal{H}$ is free as an R -module; we say that $\mathcal{H}$ is a **specialization** of $\mathcal{H}_{\mathcal{Z}}$ and call the map which sends $h \in \mathcal{H}_{\mathcal{Z}}$ to $h \otimes 1 \in \mathcal{H}$ the **specialization homomorphism**.

Let $\prime: \mathcal{Z} \rightarrow \mathcal{Z}$ be the $\mathbb{Z}$ -linear map given by $\hat{q} \mapsto \hat{q}^{-1}$ and $\hat{Q}_s \mapsto \hat{Q}_{r-s+1}$ for $1 \leq s \leq r$. Define $T'_0 = T_0$ and $T'_i = -\hat{q}^{-1}T_i$ for $1 \leq i \leq n$; using the relations of $\mathcal{H}_{\mathcal{Z}}$ it is easy to verify that $\prime$ now extends to a $\mathbb{Z}$ -linear ring involution $\prime: \mathcal{H}_{\mathcal{Z}} \rightarrow \mathcal{H}_{\mathcal{Z}}$ of $\mathcal{H}_{\mathcal{Z}}$. Hereafter, we drop the distinction between $\hat{q}$ and q , and $\hat{Q}_s$ and Q_s .

Suppose that $h \in \mathcal{H}$. Then there exists a (not necessarily unique) $h_{\mathcal{Z}} \in \mathcal{H}_{\mathcal{Z}}$ such that $h = h_{\mathcal{Z}} \otimes 1$ under specialization; we sometimes abuse notation and write

$h' = h'_Z \otimes 1 \in \mathcal{H}$. As the map $'$ does not in general define a semilinear involution on R , this notation is not well-defined on elements of $\mathcal{H}$; however, in the cases where we employ it there should be no ambiguity. For example, $T'_w = (-q)^{-\ell(w)} T_w$ and $L'_i = L_i$ for all $w \in \mathfrak{S}_n$ and $1 \leq i \leq n$.

If λ is a multipartition of n let $y_\lambda = \sum_{w \in \mathfrak{S}_\lambda} (-q)^{-\ell(w)} T_w$; then $y_\lambda = x'_\lambda$. Similarly, we define $n_\lambda = y_\lambda u_\lambda^- = m'_\lambda$ where

$$u_\lambda^- = \prod_{s=2}^r \prod_{k=1}^{\mathfrak{a}_s} (L_k - Q_{r-s+1}) = \prod_{s=1}^{r-1} \prod_{k=1}^{\mathfrak{a}_{r-s+1}} (L_k - Q_s).$$

Observe that $u_\lambda^- = (u_\lambda^+)'$; here, as usual, $\mathfrak{a}_s = |\lambda^{(1)}| + \dots + |\lambda^{(s-1)}|$ for all s . For standard tableaux $\mathfrak{s}, \mathfrak{t} \in \text{Std}(\lambda)$ set $n_{\mathfrak{s}\mathfrak{t}} = T_{d(\mathfrak{s})}^* n_\lambda T_{d(\mathfrak{t})}'$; then $m'_{\mathfrak{s}\mathfrak{t}} \in \mathcal{H}_Z$ is mapped to $n_{\mathfrak{s}\mathfrak{t}}$ under specialization. Hence, from (3.1) we obtain the following.

4.1 [13, (2.7)] *The Ariki-Koike algebra $\mathcal{H}$ is free as an R -module with cellular basis $\{n_{\mathfrak{s}\mathfrak{t}} \mid \mathfrak{s}, \mathfrak{t} \in \text{Std}(\lambda) \text{ for some } \lambda \in \Lambda^+\}$.*

Let λ be a multipartition of n . Then $(\check{\mathcal{H}}^\lambda)'$ is a two-sided ideal of $\mathcal{H}$ which is free as an R -module with basis $\{n_{\mathfrak{u}\mathfrak{v}} \mid \mathfrak{u}, \mathfrak{v} \in \text{Std}(\mu) \text{ for some } \mu \triangleright \lambda\}$. Let $\tilde{S}^\lambda$ be the Specht module (or cell module) corresponding to λ determined by the basis $\{n_{\mathfrak{s}\mathfrak{t}}\}$; then $\tilde{S}^\lambda \cong n_\lambda \mathcal{H} / (n_\lambda \mathcal{H} \cap (\check{\mathcal{H}}^\lambda)')$ and $\tilde{S}^\lambda$ is free as an R -module with basis $\{n_{\mathfrak{t}} \mid \mathfrak{t} \in \text{Std}(\lambda)\}$, where $n_{\mathfrak{t}} = n_{\mathfrak{t}\mathfrak{t}} + (\check{\mathcal{H}}^\lambda)' = (m_{\mathfrak{t}\mathfrak{t}} + \check{\mathcal{H}}^\lambda)'$ for all $\mathfrak{t} \in \text{Std}(\lambda)$.

In order to compare the two modules S^λ and $\tilde{S}^\lambda$ we need to introduce some more notation. Given a partition σ let $\sigma' = (\sigma'_1, \sigma'_2, \dots)$ be the partition which is conjugate to σ ; thus, σ'_i is the number of nodes in column i of the diagram of σ . If $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ is a multipartition then the **conjugate** $\lambda' = (\lambda'^{(1)}, \dots, \lambda'^{(r)})$ of λ' is the multipartition with $\lambda'^{(s)} = (\lambda^{(r-s+1)})'$ for $1 \leq s \leq r$.

Now suppose that $\mathfrak{t} = (\mathfrak{t}^{(1)}, \dots, \mathfrak{t}^{(r)})$ is a standard λ -tableau. Then the **conjugate** of $\mathfrak{t}$ is the standard λ' -tableau $\mathfrak{t}' = (\mathfrak{t}'^{(1)}, \dots, \mathfrak{t}'^{(r)})$ where $\mathfrak{t}'^{(s)}$ is the tableau obtained by interchanging the rows and columns of $\mathfrak{t}^{(r-s+1)}$.

With these definitions in place, we see that the following holds in Z .

4.2 *Let $\mathfrak{t}$ be a standard λ -tableau. Then $(\text{res}_{\mathfrak{t}}(k))' = \text{res}_{\mathfrak{t}'}(k)$ in Z , for $1 \leq k \leq n$.*

The expression $\text{res}_{\mathfrak{t}'}(k)$ is always well-defined; whereas $(\text{res}_{\mathfrak{t}}(k))'$ is ambiguous for certain rings R . As a first consequence we determine how the L_k act on the $n_{\mathfrak{s}\mathfrak{t}}$ -basis of $\mathcal{H}$.

4.3. Proposition. *Let $\mathfrak{s}$ and $\mathfrak{t}$ be standard λ -tableaux and suppose that k is an integer with $1 \leq m \leq n$. Then there exist $a_{\mathfrak{v}} \in R$ such that*

$$n_{\mathfrak{s}\mathfrak{t}} L_k = \text{res}_{\mathfrak{t}'}(k) n_{\mathfrak{s}\mathfrak{t}} + \sum_{\substack{\mathfrak{v} \in \text{Std}(\lambda) \\ \mathfrak{v} \triangleright \mathfrak{t}}} a_{\mathfrak{v}} n_{\mathfrak{s}\mathfrak{v}} \text{ mod } (\check{\mathcal{H}}^\lambda)'.$$

Proof. First assume that $R = Z$. Then $'$ is a $\mathbb{Z}$ -linear ring involution on $\mathcal{H}_Z$ and $L'_k = L_k$; therefore, by (3.2),

$$n_{\mathfrak{s}\mathfrak{t}} L_k = (m_{\mathfrak{s}\mathfrak{t}} L_k)' = \left(\text{res}_{\mathfrak{t}}(k) m_{\mathfrak{s}\mathfrak{t}} + \sum_{\substack{\mathfrak{v} \in \text{Std}(\lambda) \\ \mathfrak{v} \triangleright \mathfrak{t}}} a_{\mathfrak{v}} m_{\mathfrak{s}\mathfrak{v}} \text{ mod } \check{\mathcal{H}}^\lambda \right)'.$$

Using (4.2) this proves the Proposition for $\mathcal{H}_Z$. The general case now follows by specialization since $\mathcal{H} \cong \mathcal{H}_Z \otimes_Z R$. $\square$

Next consider the orthogonal basis $\{f_{\mathbf{s}\mathbf{t}}\}$ of $\mathcal{H}$ in the case where $P_{\mathcal{H}}(\mathbf{q})$ is invertible. Let $\mathcal{Z}_P$ be the localization of $\mathcal{Z}$ at $P_{\mathcal{H}}(\mathbf{q})$ and let $\mathcal{H}_{\mathcal{Z}_P}$ be the corresponding Ariki–Koike algebra. The involution $'$ extends to $\mathcal{H}_{\mathcal{Z}_P}$ and $\mathcal{H}$ is a specialization of $\mathcal{H}_{\mathcal{Z}_P}$ whenever $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R . (Note that $Q_1, \dots, Q_r$ are indeterminates in $\mathcal{Z}_P$.)

In general, $f_{\mathbf{s}\mathbf{t}} \notin \mathcal{H}_{\mathcal{Z}}$; however, if $\mathbf{t} \neq \mathbf{u}$ then $\text{res}_{\mathbf{t}}(k) - \text{res}_{\mathbf{u}}(k)$ is a factor of $P_{\mathcal{H}}(\mathbf{q})$ for all k , so $f_{\mathbf{s}\mathbf{t}} \in \mathcal{H}_{\mathcal{Z}_P}$ and we can speak of the elements $F_{\mathbf{t}}$ and $f_{\mathbf{s}\mathbf{t}} \in \mathcal{H}_{\mathcal{Z}_P}$. More generally, whenever $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R we have an element $f'_{\mathbf{s}\mathbf{t}} \in \mathcal{H}$ via specialization because $\mathcal{H} \cong \mathcal{H}_{\mathcal{Z}_P} \otimes_{\mathcal{Z}_P} R$.

4.4. Proposition. *Suppose that $\mathbf{t}$ is a standard tableau. Then $F'_{\mathbf{t}} = F_{\mathbf{t}'}$ in $\mathcal{H}_{\mathcal{Z}_P}$.*

Proof. Applying the definitions together with (4.2) gives

$$\begin{aligned} F'_{\mathbf{t}} &= \prod_{k=1}^n \prod_{\substack{c \in \mathcal{R}(k) \\ c \neq \text{res}_{\mathbf{t}}(k)}} \left(\frac{L_k - c}{\text{res}_{\mathbf{t}}(k) - c} \right)' = \prod_{k=1}^n \prod_{\substack{c \in \mathcal{R}(k) \\ c \neq \text{res}_{\mathbf{t}'}(k)}} \frac{L_k - c'}{\text{res}_{\mathbf{t}'}(k) - c'} \\ &= \prod_{k=1}^n \prod_{\substack{c \in \mathcal{R}(k) \\ c \neq \text{res}_{\mathbf{t}'}(k)}} \frac{L_k - c}{\text{res}_{\mathbf{t}'}(k) - c} = F_{\mathbf{t}'}, \end{aligned}$$

the last equality following because $\mathcal{R}(k)$ is invariant under $'$. $\square$

By Lemma 2.6 and Theorem 3.14 the Schur elements are given by $s_{\lambda}(\mathbf{q}) = \tau(F_{\mathbf{t}^{\lambda}})^{-1}$; consequently, the Schur elements have the following “palindromy” property.

4.5. Corollary. *Suppose that λ is a multipartition of n . Then $s_{\lambda'}(\mathbf{q}) = (s_{\lambda}(\mathbf{q}))'$.*

Returning to the general case, let $g_{\mathbf{s}\mathbf{t}} = F_{\mathbf{s}'} n_{\mathbf{s}\mathbf{t}} F_{\mathbf{t}'}$; then $g_{\mathbf{s}\mathbf{t}} = F'_{\mathbf{s}} m'_{\mathbf{s}\mathbf{t}} F'_{\mathbf{t}} = f'_{\mathbf{s}\mathbf{t}}$ (in $\mathcal{H}_{\mathcal{Z}_P}$). Applying $'$ to $\{f_{\mathbf{s}\mathbf{t}}\}$ and using Theorem 3.9 (and a specialization argument) shows that $\{g_{\mathbf{s}\mathbf{t}} \mid \mathbf{s}, \mathbf{t} \in \text{Std}(\lambda) \text{ for some } \lambda \in \Lambda^+\}$ is a basis of $\mathcal{H}$. Consequently, as in Corollary 3.11, $\tilde{S}^{\lambda} \cong g_{\mathbf{s}\mathbf{t}} \mathcal{H}$ for any standard λ -tableaux $\mathbf{s}, \mathbf{t} \in \text{Std}(\lambda)$.

4.6. Remark. By the Proposition and Theorem 3.14(i),

$$g_{\mathbf{t}\mathbf{t}} = f'_{\mathbf{t}\mathbf{t}} = (\gamma_{\mathbf{t}} F_{\mathbf{t}})' = \gamma'_{\mathbf{t}} F_{\mathbf{t}'} = \frac{\gamma'_{\mathbf{t}}}{\gamma_{\mathbf{t}'}} f_{\mathbf{t}'\mathbf{t}'}.$$

More generally, we can write $g_{\mathbf{s}\mathbf{t}} = \sum_{\mathbf{u}, \mathbf{v}} a_{\mathbf{u}\mathbf{v}} f_{\mathbf{u}\mathbf{v}}$ for some $a_{\mathbf{u}\mathbf{v}} \in R$. By Proposition 3.4 and Proposition 4.4, $F_{\mathbf{s}'} g_{\mathbf{s}\mathbf{t}} F_{\mathbf{t}'} = g_{\mathbf{s}\mathbf{t}}$; so it follows that $a_{\mathbf{u}\mathbf{v}} = 0$ unless $\mathbf{u} = \mathbf{s}'$ and $\mathbf{v} = \mathbf{t}'$. Therefore, $g_{\mathbf{s}\mathbf{t}} = \alpha_{\mathbf{s}\mathbf{t}} f_{\mathbf{s}'\mathbf{t}'}$ for some $\alpha_{\mathbf{s}\mathbf{t}} \in R$. Applying the $*$ -involution shows that $\alpha_{\mathbf{s}\mathbf{t}} = \alpha_{\mathbf{t}\mathbf{s}}$. Finally, by looking at the product $g_{\mathbf{s}\mathbf{t}} g_{\mathbf{t}\mathbf{s}}$ we see that $\alpha_{\mathbf{s}\mathbf{t}}^2 = \gamma'_{\mathbf{s}} \gamma'_{\mathbf{t}} / \gamma_{\mathbf{s}'} \gamma_{\mathbf{t}'}$. *A priori*, there is no reason why the square root of this element should belong to R ; nor do I see a way to determine the sign of $\alpha_{\mathbf{s}\mathbf{t}}$.

Combining Proposition 4.4 with Corollary 3.11 and the corresponding result for the g -basis shows that $S^{\lambda} \cong f_{\mathbf{t}\mathbf{t}} \mathcal{H} = g_{\mathbf{t}'\mathbf{t}'} \mathcal{H} \cong \tilde{S}^{\lambda'}$, for any $\mathbf{t} \in \text{Std}(\lambda)$. Hence, we have the following.

4.7. Corollary. *Suppose that $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R . Then $\tilde{S}^{\lambda} \cong S^{\lambda'}$.*

When R is field the assumption that $P_{\mathcal{H}}(\mathbf{q})$ is invertible is equivalent to $\mathcal{H}$ being semisimple. This assumption is necessary because, in general, $S^{\lambda'}$ and $\tilde{S}^{\lambda}$ are not isomorphic; rather, $S^{\lambda'}$ is isomorphic to the *dual* of $\tilde{S}^{\lambda}$ [24]. In the semisimple

case both S^λ and $\tilde{S}^{\lambda'}$ are irreducible, and hence self-dual, since they carry a non-degenerate bilinear form. Accordingly, we call the module $\tilde{S}^\lambda$ a **dual Specht module**.

Here is another useful application of Proposition 4.4.

4.8. Corollary. *Suppose that $\mathfrak{s}$ and $\mathfrak{t}$ are standard λ -tableaux and that $\mathfrak{u}$ and $\mathfrak{v}$ are standard μ -tableaux where λ and μ are multipartitions of n . Then $f_{\mathfrak{s}\mathfrak{t}}g_{\mathfrak{u}\mathfrak{v}} = 0$ if $\mathfrak{t} \neq \mathfrak{u}'$.*

Proof. Applying the definitions, $f_{\mathfrak{s}\mathfrak{t}}g_{\mathfrak{u}\mathfrak{v}} = F_{\mathfrak{s}}m_{\mathfrak{s}\mathfrak{t}}F_{\mathfrak{t}}F'_{\mathfrak{u}}m_{\mathfrak{u}\mathfrak{v}}F'_{\mathfrak{v}}$; however, $F'_{\mathfrak{u}} = F_{\mathfrak{u}'}$ by Proposition 4.4, so $F_{\mathfrak{t}}F'_{\mathfrak{u}} = \delta_{\mathfrak{t}\mathfrak{u}'}F_{\mathfrak{t}}$ by Theorem 3.14(i), giving the result. (By Remark 4.6, $f_{\mathfrak{s}\mathfrak{t}}g_{\mathfrak{u}'\mathfrak{v}}$ is a scalar multiple of $f_{\mathfrak{s}\mathfrak{v}'}$.) $\square$

The Specht modules S^λ and the dual Specht modules $\tilde{S}^\lambda$ are both constructed as quotient modules using the bases $\{m_{\mathfrak{s}\mathfrak{t}}\}$ and $\{n_{\mathfrak{s}\mathfrak{t}}\}$ respectively (see Corollary 3.11). In the cases where $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R we have also constructed these modules as submodules of $\mathcal{H}$. Next we produce submodules of an arbitrary Hecke algebra $\mathcal{H}$ which are isomorphic to the Specht modules; these results will also play a rôle in computing the Schur elements in the next section.

Recall that $\mathfrak{t}^\lambda$ is the λ -tableau which has the numbers $1, 2, \dots, n$ entered in order first along the rows of $\mathfrak{t}^{\lambda(1)}$ and then the rows of $\mathfrak{t}^{\lambda(2)}$ and so on. Let $\mathfrak{t}_\lambda = (\mathfrak{t}^{\lambda'})'$; that is to say that $\mathfrak{t}_\lambda$ is the λ -tableau with the numbers $1, 2, \dots, n$ entered in order first down the columns of $\mathfrak{t}_\lambda^{(r)}$ and then the columns of $\mathfrak{t}_\lambda^{(r-1)}$ etcetera. Observe that if $\mathfrak{t}$ is a standard λ -tableau then $\mathfrak{t}^\lambda \supseteq \mathfrak{t} \supseteq \mathfrak{t}_\lambda$.

4.9. Proposition. *Suppose that $P_{\mathcal{H}}(\mathbf{q})$ is invertible and let λ be a multipartition of n . Then $m_\lambda \mathcal{H} n_{\lambda'} = R f_{\mathfrak{t}^\lambda \mathfrak{t}_\lambda}$.*

Proof. By Proposition 3.4(i), $m_\lambda = f_{\mathfrak{t}^\lambda \mathfrak{t}^\lambda} + \sum_{\mathfrak{u}, \mathfrak{v} \triangleright \mathfrak{t}^\lambda} a_{\mathfrak{u}\mathfrak{v}} f_{\mathfrak{u}\mathfrak{v}}$ for some $a_{\mathfrak{u}\mathfrak{v}} \in R$. Therefore, interchanging the roles of λ and λ' and applying the involution $'$ (in $\mathcal{H}_{\mathbb{Z}_p}$ and then specializing) we see that there exist $b_{\mathfrak{a}\mathfrak{b}} \in R$ such that

$$(4.10) \quad n_{\lambda'} = g_{\mathfrak{t}^{\lambda'} \mathfrak{t}^{\lambda'}} + \sum_{\mathfrak{a}, \mathfrak{b} \triangleright \mathfrak{t}^{\lambda'}} b_{\mathfrak{a}\mathfrak{b}} g_{\mathfrak{a}\mathfrak{b}} = \frac{\gamma'_{\mathfrak{t}^{\lambda'}}}{\gamma_{\mathfrak{t}^\lambda}} f_{\mathfrak{t}^\lambda \mathfrak{t}_\lambda} + \sum_{\mathfrak{t}_\lambda \triangleright \mathfrak{a}', \mathfrak{b}'} b_{\mathfrak{a}\mathfrak{b}} g_{\mathfrak{a}\mathfrak{b}}$$

where for the second equality we have used Remark 4.6 (note that $(\mathfrak{t}^{\lambda'})' = \mathfrak{t}_\lambda$) and the observation that $\mathfrak{a}, \mathfrak{b} \triangleright \mathfrak{t}^{\lambda'}$ if and only if $\mathfrak{t}_\lambda \triangleright \mathfrak{a}', \mathfrak{b}'$. Now $m_\lambda \mathcal{H} n_{\lambda'}$ is spanned by the elements $m_\lambda f_{\mathfrak{s}\mathfrak{t}} n_{\lambda'}$, where $\mathfrak{s}$ and $\mathfrak{t}$ range over all pairs of standard tableaux of the same shape. Now, (4.10) and Corollary 4.8 imply that

$$\begin{aligned} m_\lambda f_{\mathfrak{s}\mathfrak{t}} n_{\lambda'} &= \left(f_{\mathfrak{t}^\lambda \mathfrak{t}^\lambda} + \sum_{\mathfrak{u}, \mathfrak{v} \triangleright \mathfrak{t}^\lambda} a_{\mathfrak{u}\mathfrak{v}} f_{\mathfrak{u}\mathfrak{v}} \right) f_{\mathfrak{s}\mathfrak{t}} \left(\frac{\gamma'_{\mathfrak{t}^{\lambda'}}}{\gamma_{\mathfrak{t}^\lambda}} f_{\mathfrak{t}_\lambda \mathfrak{t}_\lambda} + \sum_{\mathfrak{t}_\lambda \triangleright \mathfrak{a}', \mathfrak{b}'} b_{\mathfrak{a}\mathfrak{b}} g_{\mathfrak{a}\mathfrak{b}} \right) \\ &= \frac{\gamma'_{\mathfrak{t}^{\lambda'}}}{\gamma_{\mathfrak{t}^\lambda}} f_{\mathfrak{t}^\lambda \mathfrak{t}^\lambda} f_{\mathfrak{s}\mathfrak{t}} f_{\mathfrak{t}_\lambda \mathfrak{t}_\lambda} + \sum_{\substack{\mathfrak{t}_\lambda \triangleright \mathfrak{a}', \mathfrak{b}' \\ \mathfrak{u}, \mathfrak{v} \triangleright \mathfrak{t}^\lambda}} a_{\mathfrak{u}\mathfrak{v}} b_{\mathfrak{a}\mathfrak{b}} f_{\mathfrak{u}\mathfrak{v}} f_{\mathfrak{s}\mathfrak{t}} g_{\mathfrak{a}\mathfrak{b}}, \\ &= \frac{\gamma'_{\mathfrak{t}^{\lambda'}}}{\gamma_{\mathfrak{t}^\lambda}} f_{\mathfrak{t}^\lambda \mathfrak{t}^\lambda} f_{\mathfrak{s}\mathfrak{t}} f_{\mathfrak{t}_\lambda \mathfrak{t}_\lambda} \\ &= \begin{cases} \gamma_{\mathfrak{t}^\lambda} \gamma'_{\mathfrak{t}^{\lambda'}} f_{\mathfrak{t}^\lambda \mathfrak{t}_\lambda}, & \text{if } \mathfrak{s} = \mathfrak{t}^\lambda \text{ and } \mathfrak{t} = \mathfrak{t}_\lambda, \\ 0, & \text{otherwise,} \end{cases} \end{aligned}$$

with the last equality following from Theorem 3.9. Therefore, $m_\lambda \mathcal{H} n_{\lambda'} = R f_{\mathfrak{t}^\lambda \mathfrak{t}_\lambda}$ as required. $\square$

Let $w_\lambda = d(\mathbf{t}_\lambda)$; thus, w_λ is the unique element of $\mathfrak{S}_n$ such that $\mathbf{t}_\lambda = \mathbf{t}^\lambda w_\lambda$.

4.11. *Example.* Let $\lambda = ((2, 1^2), (2, 1), (2))$. Then

$$\mathbf{t}^\lambda = \left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline 4 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 5 & 6 \\ \hline 7 & \\ \hline & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 8 & 9 \\ \hline & \\ \hline & \\ \hline \end{array} \right) \quad \text{and} \quad \mathbf{t}_\lambda = \left(\begin{array}{|c|c|} \hline 6 & 9 \\ \hline 7 & \\ \hline 8 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & 5 \\ \hline 4 & \\ \hline & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline & \\ \hline & \\ \hline \end{array} \right)$$

and $w_\lambda = (1, 6, 5, 3, 7, 4, 8)(2, 9)$.

In order to compute the Schur elements we will need to know quite a few properties of the permutations w_λ ; these permutations enter the story through the following definition and Corollary.

4.12. Definition. Suppose that λ is a multipartition of n . Let $z_\lambda = m_\lambda T_{w_\lambda} n_{\lambda'}$.

The element z_λ and the following result are crucial to our computation of the Schur elements.

4.13. Corollary. Suppose that $P_{\mathcal{H}}(\mathbf{q})$ is invertible in R . Then $z_\lambda = \gamma'_{\mathbf{t}^\lambda} f_{\mathbf{t}^\lambda \mathbf{t}_\lambda}$. In particular, $m_\lambda \mathcal{H} n_{\lambda'} = R z_\lambda$.

Proof. Consulting the definitions, $z_\lambda = m_\lambda T_{w_\lambda} n_{\lambda'} = m_{\mathbf{t}^\lambda \mathbf{t}_\lambda} n_{\lambda'}$. Furthermore, by Proposition 3.4(i), there exist $c_{\mathbf{u}\mathbf{v}} \in R$ such that $m_{\mathbf{t}^\lambda \mathbf{t}_\lambda} = f_{\mathbf{t}^\lambda \mathbf{t}_\lambda} + \sum c_{\mathbf{u}\mathbf{v}} f_{\mathbf{u}\mathbf{v}}$ where the sum is over the pairs $(\mathbf{u}, \mathbf{v})$ of standard tableaux which strictly dominate $\mathbf{t}_\lambda$. Therefore, by (4.10),

$$z_\lambda = \left(f_{\mathbf{t}^\lambda \mathbf{t}_\lambda} + \sum_{\mathbf{u}, \mathbf{v} \triangleright \mathbf{t}_\lambda} c_{\mathbf{u}\mathbf{v}} f_{\mathbf{u}\mathbf{v}} \right) \left(\frac{\gamma'_{\mathbf{t}^\lambda}}{\gamma_{\mathbf{t}_\lambda}} f_{\mathbf{t}_\lambda \mathbf{t}_\lambda} + \sum_{\substack{\mathbf{a}, \mathbf{b} \\ \mathbf{t}_\lambda \triangleright \mathbf{a}', \mathbf{b}'}} b_{\mathbf{a}\mathbf{b}} g_{\mathbf{a}\mathbf{b}} \right).$$

By Theorem 3.9, $f_{\mathbf{t}^\lambda \mathbf{t}_\lambda} f_{\mathbf{t}_\lambda \mathbf{t}_\lambda} = \gamma_{\mathbf{t}_\lambda} f_{\mathbf{t}^\lambda \mathbf{t}_\lambda}$ and this is the only non-zero term in this product by Corollary 4.8. Hence, $z_\lambda = \gamma'_{\mathbf{t}^\lambda} f_{\mathbf{t}^\lambda \mathbf{t}_\lambda}$ as required. $\square$

Now, z_λ is an element of $\mathcal{H}_{\mathbf{Z}}$, so $\gamma'_{\mathbf{t}^\lambda} f_{\mathbf{t}^\lambda \mathbf{t}_\lambda} \in \mathcal{H}_{\mathbf{Z}}$. By definition, $z_\lambda \mathcal{H}$ is a submodule of $m_\lambda \mathcal{H}$ and a quotient module of $n_{\lambda'} \mathcal{H}$. Over an arbitrary ring R , Du and Rui [13, Remark 2.5] showed that $S^\lambda \cong z_\lambda^* \mathcal{H}$ and $\tilde{S}^{\lambda'} \cong z_\lambda \mathcal{H}$ as $\mathcal{H}$ -modules, the isomorphisms being given by the natural quotient maps $m_\lambda \mathcal{H} \rightarrow z_\lambda^* \mathcal{H}$ and $n_{\lambda'} \mathcal{H} \rightarrow z_\lambda \mathcal{H}$. Note that $S^\lambda \cong \tilde{S}^{\lambda'}$ when $\mathcal{H}$ is semisimple by Corollary 3.11.

5. THE SCHUR ELEMENTS

Using the results of the previous sections we are now ready to compute the Schur elements of $\mathcal{H}$. By Lemma 2.6, $s_\lambda(\mathbf{q}) = \frac{1}{\tau(F_{\mathbf{t}^\lambda})}$, so it is enough to calculate $\tau(F_{\mathbf{t}^\lambda})$. Our basic strategy, which is inspired by Murphy [25], is to write $F_{\mathbf{t}^\lambda}$ as a product of two terms and, in effect, to evaluate τ on each of these factors separately.

5.1. Proposition. Suppose that $\mathbf{t}$ is a standard λ -tableau. Then there exist elements $\Phi_{\mathbf{t}}$ and $\Psi_{\mathbf{t}}$ in $\mathcal{H}(\mathfrak{S}_n)$ such that

- (i) $\Psi_{\mathbf{t}} F_{\mathbf{t}} = F_{\mathbf{t}^\lambda} \Phi_{\mathbf{t}}$;
- (ii) $\Psi_{\mathbf{t}} = T_{d(\mathbf{t})} + \sum_{w < d(\mathbf{t})} p_{\mathbf{t}w} T_w$, for some $p_{\mathbf{t}w} \in R$; and,
- (iii) $\gamma_{\mathbf{t}^\lambda} \Phi_{\mathbf{t}} \Psi_{\mathbf{t}}^* = \gamma_{\mathbf{t}}$.

Proof. We prove all three statements by induction on $\mathbf{t}$. When $\mathbf{t} = \mathbf{t}^\lambda$ there is nothing to prove as we may take $\Phi_{\mathbf{t}^\lambda} = \Psi_{\mathbf{t}^\lambda} = 1$. Suppose then that $\mathbf{t} \neq \mathbf{t}^\lambda$. Then there exists an integer i , with $1 \leq i < n$, such that $\mathbf{s} = \mathbf{t}(i, i+1) \triangleright \mathbf{t}$. Let $\alpha = \frac{(q-1)\text{res}_{\mathbf{t}}(i)}{\text{res}_{\mathbf{t}}(i) - \text{res}_{\mathbf{s}}(i)}$ and $\beta = \frac{(q-1)\text{res}_{\mathbf{s}}(i)}{\text{res}_{\mathbf{s}}(i) - \text{res}_{\mathbf{t}}(i)}$. Then $f_{\mathbf{s}\mathbf{s}}(T_i - \alpha) = f_{\mathbf{s}\mathbf{t}}$ by Proposition 3.5. Similarly, by the left hand analogue of Proposition 3.5 (interchanging the roles of $\mathbf{s}$ and $\mathbf{t}$), together with (3.7)(ii), $(T_i - \beta)f_{\mathbf{t}\mathbf{t}} = (\gamma_{\mathbf{t}}/\gamma_{\mathbf{s}})f_{\mathbf{s}\mathbf{t}}$. Therefore,

$$(T_i - \beta)F_{\mathbf{t}} = \frac{1}{\gamma_{\mathbf{t}}}(T_i - \beta)f_{\mathbf{t}\mathbf{t}} = \frac{1}{\gamma_{\mathbf{s}}}f_{\mathbf{s}\mathbf{t}} = \frac{1}{\gamma_{\mathbf{s}}}f_{\mathbf{s}\mathbf{s}}(T_i - \alpha) = F_{\mathbf{s}}(T_i - \alpha).$$

By induction, there exist elements $\Phi_{\mathbf{s}}$ and $\Psi_{\mathbf{s}}$ which satisfy properties (i)–(iii). Define $\Psi_{\mathbf{t}} = \Psi_{\mathbf{s}}(T_i - \beta)$ and $\Phi_{\mathbf{t}} = \Phi_{\mathbf{s}}(T_i - \alpha)$; then, by induction and the last equation,

$$\Psi_{\mathbf{t}}F_{\mathbf{t}} = \Psi_{\mathbf{s}}(T_i - \beta)F_{\mathbf{t}} = \Psi_{\mathbf{s}}F_{\mathbf{s}}(T_i - \alpha) = F_{\mathbf{t}^\lambda}\Phi_{\mathbf{s}}(T_i - \alpha) = F_{\mathbf{t}^\lambda}\Phi_{\mathbf{t}}.$$

Hence, (i) holds. Next, again by induction we have

$$\Psi_{\mathbf{t}} = \Psi_{\mathbf{s}}(T_i - \beta) = \left(T_{d(\mathbf{s})} + \sum_{v < d(\mathbf{s})} p_{sv}T_v\right)(T_i - \beta) = T_{d(\mathbf{t})} + \sum_{w < d(\mathbf{t})} p_{tw}T_w,$$

by standard properties of the Bruhat order since $d(\mathbf{t}) = d(\mathbf{s})(i, i+1) > d(\mathbf{s})$. This proves (ii). Finally, using induction once more (and a quick calculation for the second equality),

$$\begin{aligned} \gamma_{\mathbf{t}^\lambda}\Phi_{\mathbf{t}}\Psi_{\mathbf{t}}^* &= \gamma_{\mathbf{t}^\lambda}\Phi_{\mathbf{s}}(T_i - \alpha)(T_i - \beta)\Psi_{\mathbf{s}}^* \\ &= \frac{(q\text{res}_{\mathbf{s}}(i) - \text{res}_{\mathbf{t}}(i))(\text{res}_{\mathbf{s}}(i) - q\text{res}_{\mathbf{t}}(i))}{(\text{res}_{\mathbf{s}}(i) - \text{res}_{\mathbf{t}}(i))^2}\gamma_{\mathbf{t}^\lambda}\Phi_{\mathbf{s}}\Psi_{\mathbf{s}}^* \\ &= \frac{(q\text{res}_{\mathbf{s}}(i) - \text{res}_{\mathbf{t}}(i))(\text{res}_{\mathbf{s}}(i) - q\text{res}_{\mathbf{t}}(i))}{(\text{res}_{\mathbf{s}}(i) - \text{res}_{\mathbf{t}}(i))^2}\gamma_{\mathbf{s}} = \gamma_{\mathbf{t}}, \end{aligned}$$

the last equality coming from (3.7)(ii). This proves (iii) and so completes the proof. $\square$

We are not claiming that the elements $\Phi_{\mathbf{t}}$ and $\Psi_{\mathbf{t}}$ are uniquely determined by the conditions of the Proposition; ostensibly, these elements depend upon the choice of reduced expression for $d(\mathbf{t})$. In what follows we only need to know that elements with these properties exist.

5.2. Corollary. *Suppose that $\mathbf{t}$ is a standard λ -tableau. Then*

- (i) $F_{\mathbf{t}^\lambda} = \frac{\gamma_{\mathbf{t}^\lambda}}{\gamma_{\mathbf{t}}}\Psi_{\mathbf{t}}F_{\mathbf{t}}\Psi_{\mathbf{t}}^*$; and,
- (ii) $F_{\mathbf{t}} = \frac{\gamma_{\mathbf{t}^\lambda}}{\gamma_{\mathbf{t}}}\Phi_{\mathbf{t}}^*F_{\mathbf{t}^\lambda}\Phi_{\mathbf{t}}$.

Proof. Using parts (iii) and (i) of the Proposition, respectively, shows that

$$F_{\mathbf{t}^\lambda} = \frac{\gamma_{\mathbf{t}^\lambda}}{\gamma_{\mathbf{t}}}F_{\mathbf{t}^\lambda}\Phi_{\mathbf{t}}\Psi_{\mathbf{t}}^* = \frac{\gamma_{\mathbf{t}^\lambda}}{\gamma_{\mathbf{t}}}\Psi_{\mathbf{t}}F_{\mathbf{t}}\Psi_{\mathbf{t}}^*;$$

this proves (i). Part (ii) follows from (i) by ‘conjugating’ (i) by $\Phi_{\mathbf{t}}$. $\square$

The main reason why we are interested in $\Psi_{\mathbf{t}}$ and $\Phi_{\mathbf{t}}$ is the following.

5.3. Proposition. *Suppose that $\mathbf{s}$ and $\mathbf{t}$ are standard λ -tableaux. Then*

$$f_{\mathbf{s}\mathbf{t}} = \Phi_{\mathbf{s}}^*f_{\mathbf{t}^\lambda\mathbf{t}}\Phi_{\mathbf{t}}.$$

Proof. By the definition of f_{st} and Proposition 5.1(ii) we have

$$f_{st} = F_s T_{d(s)}^* m_\lambda T_{d(t)} F_t = F_s \Psi_s^* m_\lambda \Psi_t F_t - \sum_{(v,w) < (d(s), d(t))} p_{sv} p_{tw} F_s T_v^* m_\lambda T_w F_t.$$

Now if $(v, w) < (d(s), d(t))$ then $T_v^* m_\lambda T_w$ belongs to the span of the m_{uv} where $(u, v) \triangleright (s, t)$. Therefore, by Proposition 3.4(i), $T_v^* m_\lambda T_w$ belongs to the span of the f_{uv} where either u and v are standard λ -tableaux and $(u, v) \triangleright (s, t)$, or $\text{Shape}(u) = \text{Shape}(v) \triangleright \lambda$; consequently, $F_s T_v^* m_\lambda T_w F_t = 0$ by Proposition 3.4(iii). Hence, by Theorem 3.14(i) and Proposition 5.1(i),

$$f_{st} = F_s \Psi_s^* m_\lambda \Psi_t F_t = \Phi_s^* F_{t^\lambda} m_\lambda F_{t^\lambda} \Phi_t = \Phi_s^* f_{t^\lambda t^\lambda} \Phi_t$$

as required. $\square$

Applying this result to $f_{t^\lambda t^\lambda}$ shows that $f_{t^\lambda t^\lambda} = \Phi_{t^\lambda}^* f_{t^\lambda t^\lambda} \Phi_{t^\lambda} = \gamma_{t^\lambda} F_{t^\lambda} \Phi_{t^\lambda}$, the last equality following because $\Phi_{t^\lambda} = 1$. Using Proposition 5.1(iii) to multiply this equation on the right by $\Psi_{t^\lambda}^*$, and recalling Corollary 4.13, now yields the following.

5.4. Corollary. *Suppose that λ is a multipartition of n . Then*

$$F_{t^\lambda} = \frac{1}{\gamma_{t^\lambda}} f_{t^\lambda t^\lambda} \Psi_{t^\lambda}^* = \frac{1}{\gamma_{t^\lambda} \gamma_{t^\lambda}'} z_\lambda \Psi_{t^\lambda}^*.$$

By Lemma 2.6 in order to compute the Schur elements it suffices to calculate $\tau(F_{t^\lambda})$ for each λ ; so we are reduced to finding $\tau(z_\lambda \Psi_{t^\lambda}^*)$. To do this we rewrite z_λ with respect to the Ariki–Koike basis. In types A and B (that is, $r = 1$ or $r = 2$) this is reasonably straightforward; in general we have to work much harder.

Until further notice, fix a multipartition λ and let $\bar{\lambda} = (\bar{\lambda}^{(1)}, \dots, \bar{\lambda}^{(r)})$ be the multipartition with $\bar{\lambda}^{(s)} = (|\lambda^{(s)}|)$ for $1 \leq s \leq r$. Then $w_{\bar{\lambda}} = d(t^{\bar{\lambda}})$ is the permutation

$$\left(\begin{array}{ccc|ccc|ccc} 1 & \cdots & a_2 & a_2 + 1 & \cdots & a_3 & \cdots & a_r + 1 & \cdots & n \\ b_1 + 1 & \cdots & n & b_2 + 1 & \cdots & b_1 & \cdots & 1 & \cdots & b_{r-1} \end{array} \right),$$

where $a_s = \sum_{t=1}^{s-1} |\lambda^{(t)}|$ and $b_s = \sum_{t=s+1}^r |\lambda^{(t)}|$ for $1 \leq s \leq r$.

Now, $t^{\bar{\lambda}} w_{\bar{\lambda}}$ is a standard tableau; therefore, $w_{\bar{\lambda}}$ is a distinguished right coset representative for $\mathfrak{S}_{\bar{\lambda}}$ in $\mathfrak{S}_n$ (for example, by [22, Prop. 3.3]). Equivalently, $w_{\bar{\lambda}}$ is a distinguished left coset representative for $\mathfrak{S}_{\bar{\lambda}}$ in $\mathfrak{S}_n$ since $\mathfrak{S}_{\bar{\lambda}} w_{\bar{\lambda}} = w_{\bar{\lambda}} \mathfrak{S}_{\bar{\lambda}}$. Consequently, there exists $w_{\lambda/\bar{\lambda}} \in \mathfrak{S}_{\bar{\lambda}}$ such that $w_\lambda = w_{\bar{\lambda}} w_{\lambda/\bar{\lambda}}$ and $\ell(w_\lambda) = \ell(w_{\bar{\lambda}}) + \ell(w_{\lambda/\bar{\lambda}})$. To proceed we need to factorize $w_{\bar{\lambda}}$.

Suppose that a and b are non-negative integers let $w_{a,b} = w_{((a),(b))}$; so, $t_{((a),(b))} = t^{((a),(b))} w_{a,b}$. More concretely, $w_{a,0} = 1 = w_{0,b}$ and if $a > 0$ and $b > 0$ then

$$w_{a,b} = \left(\begin{array}{cccc|cccc} 1 & 2 & \cdots & a & a+1 & a+2 & \cdots & a+b \\ b+1 & b+2 & \cdots & a+b & 1 & 2 & \cdots & b \end{array} \right).$$

If $i \leq j$ we also set $s_{i,j} = s_i s_{i+1} \dots s_j$ and $s_{j,i} = 1$; so $s_{i,j} = (j+1, j, \dots, i)$. For convenience let $T_{i,j} = T_{s_{i,j}}$. It is not hard to see that $w_{a,b} = (s_{a+b-1,1})^b$.

The permutations $w_{a,b}$ were studied by Dipper and James [11]; in particular, they observed that $w_{a,b} = s_{a,a+b-1} w_{a-1,b}$, with the lengths adding, which implies the following. (Note that our $s_{i,j}$ is Dipper and James' $s_{i,j-1}$ when $i < j$; also if $i \geq j$ then Dipper and James set $s_{i,j} = (s_{j,i-1})^{-1}$ whereas we have $s_{i,j} = 1$.) We let $\mathfrak{S}_{(b,a)} = \mathfrak{S}_b \times \mathfrak{S}_a \hookrightarrow \mathfrak{S}_n$ (natural embedding).

5.5. Lemma. *Suppose that a and b are positive integers with $a + b \leq n$. Then $w_{a,b} = s_{a,a+b-1} \dots s_{1,b}$ and $\ell(w_{a,b}) = \ell(s_{a,a+b-1}) + \dots + \ell(s_{1,b})$. Moreover, $w_{a,b}$ is a distinguished left coset representative for $\mathfrak{S}_{(b,a)}$ in $\mathfrak{S}_n$; that is, $\ell(w_{a,b}v) = \ell(w_{a,b}) + \ell(v)$ for all $v \in \mathfrak{S}_{(b,a)}$.*

For $s = 1, \dots, r$ define $w_{\bar{\lambda},s} = w_{n_s, \mathfrak{b}_s}$, where $n_s = |\lambda^{(s)}|$ and $\mathfrak{b}_s = \sum_{t=s+1}^r |\lambda^{(t)}|$; in particular, $\mathfrak{b}_0 = n$ and $w_{\bar{\lambda},r} = 1$. Now $n_s + \mathfrak{b}_s = \mathfrak{b}_{s-1}$ so $w_{\bar{\lambda},s} \in \mathfrak{S}_{\mathfrak{b}_{s-1}}$ for each s ; therefore, $w_{\bar{\lambda},s} \dots w_{\bar{\lambda},r-1}$ is an element of $\mathfrak{S}_{\mathfrak{b}_{s-1}}$ and, consequently,

$$\ell(w_{\bar{\lambda},1} \dots w_{\bar{\lambda},r-1}) = \ell(w_{\bar{\lambda},1}) + \ell(w_{\bar{\lambda},2} \dots w_{\bar{\lambda},r-1}) = \dots = \ell(w_{\bar{\lambda},1}) + \dots + \ell(w_{\bar{\lambda},r-1})$$

by Lemma 5.5. Noting that $(k)w_{\bar{\lambda}} = (k)w_{\bar{\lambda},1} \dots w_{\bar{\lambda},s}$ for $k = 1, 2, \dots, \mathfrak{a}_{s+1}$ we have shown the following.

5.6 *Let λ be a partition of n . Then $w_{\lambda} = w_{\bar{\lambda}}w_{\lambda/\bar{\lambda}} = w_{\bar{\lambda},1} \dots w_{\bar{\lambda},r-1}w_{\lambda/\bar{\lambda}}$. Moreover, $\ell(w_{\lambda}) = \ell(w_{\bar{\lambda}}) + \ell(w_{\lambda/\bar{\lambda}}) = \ell(w_{\bar{\lambda},1}) + \dots + \ell(w_{\bar{\lambda},r-1}) + \ell(w_{\lambda/\bar{\lambda}})$.*

The point of the lengths adding is that $T_{w_{\lambda}} = T_{w_{\bar{\lambda},1}} \dots T_{w_{\bar{\lambda},r-1}}T_{w_{\lambda/\bar{\lambda}}}$. We will use this below without further comment.

Although we won't need it notice that $\ell(w_{a,n-a}) = a(n-a)$ by Lemma 5.5 (or directly); so (5.6) implies that $\ell(w_{\bar{\lambda}}) = \sum_{1 \leq s < t \leq r} |\lambda^{(s)}||\lambda^{(t)}|$.

5.7. Example. Let $\lambda = ((2, 1^2), (2, 1), (2))$. Then $\bar{\lambda} = ((4), (3), (2))$ and

$$\begin{aligned} w_{\lambda} &= \left(\begin{array}{cccc|cccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 6 & 9 & 7 & 8 & 3 & 5 & 4 & 1 \end{array} \begin{array}{cc} 9 & 2 \end{array} \right) = (1, 6, 5, 3, 7, 4, 8)(2, 9), \\ w_{\bar{\lambda}} &= \left(\begin{array}{cccc|cccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 6 & 7 & 8 & 9 & 3 & 4 & 5 & 1 \end{array} \begin{array}{cc} 9 & 2 \end{array} \right) = (1, 6, 4, 9, 2, 7, 5, 3, 8); \end{aligned}$$

therefore, $w_{\lambda/\bar{\lambda}} = (7, 9, 8)(4, 5)$. Further, $\mathfrak{b} = (5, 2, 0)$ so, using Lemma 5.5,

$$\begin{aligned} w_{\bar{\lambda},1} &= w_{4,5} = \left(\begin{array}{cccc|cccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 6 & 7 & 8 & 9 & 1 & 2 & 3 & 4 \end{array} \begin{array}{cc} 9 & 5 \end{array} \right) \\ &= (1, 6, 2, 7, 3, 8, 4, 9, 5) = s_{4,8}s_{3,7}s_{2,6}s_{1,5} \\ w_{\bar{\lambda},2} &= w_{3,2} = \left(\begin{array}{ccc|cc} 1 & 2 & 3 & 4 & 5 \\ 3 & 4 & 5 & 1 & 2 \end{array} \right) = (1, 3, 5, 2, 4) = s_{3,4}s_{2,3}s_{1,2} \end{aligned}$$

and $w_{\bar{\lambda},3} = 1$. It is evident that $w_{\lambda} = w_{\bar{\lambda},1}w_{\bar{\lambda},2}w_{\lambda/\bar{\lambda}}$.

Given $i \leq j$ and $Q \in R$ let $L_{i,j}(Q) = (L_i - Q) \dots (L_j - Q)$; if $i > j$ set $L_{i,j}(Q) = 1$. In particular, $u_{\bar{\lambda}}^- = L_{1,\mathfrak{b}_1}(Q_1) \dots L_{1,\mathfrak{b}_{r-1}}(Q_{r-1})$ where $\mathfrak{b}_s = |\lambda^{(s+1)}| + \dots + |\lambda^{(r)}|$, as above.

As a final piece of notation, given $0 \leq l \leq m < n$ let $H_{k,m}$ be the R -submodule of $\mathcal{H}$ spanned by the elements $\{L_1^{c_1} \dots L_k^{c_k} T_w \mid 0 \leq c_i < r \text{ and } w \in \mathfrak{S}_m\}$. In general $H_{k,m}$ is neither a subalgebra nor a submodule of $\mathcal{H}$; however, $H_{k,m}$ is a right $\mathcal{H}(\mathfrak{S}_m)$ -module.

One of the difficulties in working with $\mathcal{H}$ is that, in general, the powers of the L_k are not necessarily linear combinations of products of the L_i^s , for $1 \leq i \leq k$ and $1 \leq s < r$; however, it is always true that $L_k^s \in H_{k,m}$ for all $s \geq 0$ provided that $m \geq k$ (since in this case $\langle T_0, \dots, T_{k-1} \rangle \subseteq H_{k,m}$ by (2.2)).

5.8. Lemma. *Suppose that $h \in H_{k-1,k}$ for some k with $1 \leq k \leq n$.*

(i) If $1 \leq i < n$ then

$$T_i h \in \begin{cases} H_{k-1,k}, & \text{if } 1 \leq i < k-1, \\ H_{k,k}, & \text{if } i = k-1, \\ H_{k-1,i+1}, & \text{if } k \leq i < n. \end{cases}$$

(ii) If $1 \leq s < r$ then $hL_k^s = \sum_{t=0}^s L_k^t h_t$ for some $h_t \in H_{k-1,k}$.

Proof. By definition h is a linear combination of terms of the form $L_{k-1}^{c_{k-1}} \dots L_1^{c_1} T_w$, for some $0 \leq c_i < r$ and $w \in \mathfrak{S}_k$. By (2.1)(ii), if $i \neq m-1, m$ then $T_i L_m = L_m T_i$; whereas $T_m L_m = L_{m+1}(T_m - q + 1)$ and $T_{m-1} L_m = L_{m-1} T_{m-1} + (q-1)L_m$. Combining these facts proves (i).

For part (ii), observe that if $w \in \mathfrak{S}_k$ then $T_w L_k = L_k T_w$ unless s_{k-1} appears in a reduced expression of w ; however, $T_{k-1} L_k = L_{k-1} T_{k-1} + (q-1)L_k$. Part (ii) now follows by induction on s using the remarks of the last paragraph. $\square$

5.9. Lemma. Fix $Q \in R$ and let i, j, c and d be positive integers with $i \leq c \leq d \leq j < n$. Then

$$T_{i,j} L_{c,d}(Q) = \left\{ L_{c+1,d+1}(Q) + \sum_{k=c+1}^{d+1} L_k h_k \right\} T_{i,j}$$

for some elements $h_k \in H_{k-1,k}$ for $c+1 \leq k \leq d+1$.

Proof. If $d = j$ then by convention $T_{d+1,j} = 1$. Therefore, by (2.1)(iv) we have

$$\begin{aligned} T_{i,j} L_{c,d}(Q) &= T_{i,c-1} T_{c,d}(L_c - Q) \dots (L_d - Q) T_{d+1,j} \\ &= T_{i,c-1} T_c(L_c - Q) \dots T_d(L_d - Q) T_{d+1,j}. \end{aligned}$$

Now, $T_k L_k = q L_{k+1} T_k^{-1} = L_{k+1} T_k - (q-1)L_{k+1}$, so

$$T_{i,j} L_{c,d}(Q) = T_{i,c-1} \prod_{k=c+1}^{d+1} \left\{ (L_k - Q) T_{k-1} - (q-1)L_k \right\} \cdot T_{d+1,j},$$

where we read the terms from left to right with increasing values of k — the order of the factors is important here. Now T_{k-1} commutes with L_m if $k < m$; therefore, each T_{k-1} in this product commutes with all of L_m 's which appear to its right. A straightforward induction on $(d-c)$ using (2.1) shows that there exist $h'_k \in H_{k-1,k}$ such that

$$T_{i,j} L_{c,d}(Q) = T_{i,c-1} \left\{ L_{c+1,d+1}(Q) T_{c,d} + \sum_{k=c+1}^{d+1} L_k h'_k T_{k,d} \right\} T_{d+1,j}.$$

(Recall that our convention is that $T_{d+1,d} = 1$.) If $c+1 \leq k \leq d+1$ then by (2.1)

$$T_{i,c-1} L_k h'_k T_{k,d} T_{d+1,j} = L_k T_{i,c-1} h'_k T_{k,j} = L_k T_{i,c-1} h'_k T_{i,k-1}^{-1} T_{i,j}.$$

Also, using (2.1)(ii), $T_{i,c-1} L_{c+1,d+1}(Q) = L_{c+1,d+1}(Q) T_{i,c-1}$. Therefore,

$$T_{i,j} L_{c,d}(Q) = L_{c+1,d+1}(Q) T_{i,j} + \sum_{k=c+1}^{d+1} L_k h_k T_{i,j},$$

where $h_k = T_{i,c-1} h'_k T_{i,k-1}^{-1} \in H_{k-1,k}$ by Lemma 5.8, for $c+1 \leq k \leq d+1$. $\square$

This brings us to the key technical lemma.

5.10. Lemma. *Fix $Q \in R$ and let a, b, c and d be non-negative integers such that $a + b \leq n$ and $1 \leq c \leq d \leq b$. Then there exist elements $h_k \in H_{k-1,k}$ such that*

$$T_{w_{a,b}} L_{c,d}(Q) = \left\{ L_{a+c,a+d}(Q) + \sum_{k=a+c}^{a+d} L_k h_k \right\} T_{w_{a,b}}.$$

Proof. We argue by induction on a . If $a = 0$ then $w_{a,b} = 1$ and there is nothing to prove; so suppose that $a \geq 1$. Then $w_{a,b} = s_{a,a+b-1} w_{a-1,b}$ by Lemma 5.5. Therefore, by induction and Lemma 5.9, respectively, there exist elements $h'_k, h''_k \in H_{k-1,k}$ such that

$$\begin{aligned} T_{w_{a,b}} L_{c,d}(Q) &= T_{a,a+b-1} T_{w_{a-1,b}} L_{c,d}(Q) \\ &= T_{a,a+b-1} \left\{ L_{a+c-1,a+d-1}(Q) + \sum_{k=a+c-1}^{a+d-1} L_k h'_k \right\} T_{w_{a-1,b}} \\ &= \left\{ L_{a+c,a+d}(Q) + \sum_{k=a+c}^{a+d} L_k h''_k \right\} T_{a,a+b-1} T_{w_{a-1,b}} \\ &\quad + \sum_{k=a+c-1}^{a+d-1} T_{a,a+b-1} L_k h'_k T_{w_{a-1,b}}. \end{aligned}$$

If $a + c - 1 \leq k \leq a + d - 1$ then by (2.1)

$$\begin{aligned} T_{a,a+b-1} L_k h'_k T_{w_{a-1,b}} &= T_{a,k-1} T_k L_k T_{k+1,a+b-1} h'_k T_{w_{a-1,b}} \\ &= q T_{a,k-1} L_{k+1} T_k^{-1} T_{k+1,a+b-1} h'_k T_{w_{a-1,b}} \\ &= q L_{k+1} T_{a,k-1} T_k^{-1} h'_k T_{k+1,a+b-1} T_{w_{a-1,b}} \\ &= q L_{k+1} T_{a,k-1} T_k^{-1} h'_k T_{a,k}^{-1} T_{a,a+b-1} T_{w_{a-1,b}} \\ &= L_{k+1} T_{a,k-1} (T_k - q + 1) h'_k T_{a,k}^{-1} T_{w_{a,b}}, \end{aligned}$$

where the third equality follows by (2.1) because $h'_k \in H_{k-1,k}$. By Lemma 5.8 the element $T_{a,k-1} (T_k - q + 1) h'_k T_{a,k}^{-1}$ belongs to $H_{k,k+1}$; so, combining these equations proves the Lemma. $\square$

Lemma 5.10 is enough to compute the Schur elements when $r = 2$; in order to cover the general case we delicately apply Lemma 5.10 several times.

5.11. Corollary. *Suppose that a, b, c and s are non-negative integers with $a + b \leq n$, $1 \leq c \leq b$ and $1 \leq s < r$. Then $T_{w_{a,b}} L_c^s = \sum_{t=1}^s L_{a+c}^t h_t T_{w_{a,b}}$ some $h_t \in H_{a+c-1,a+c}$.*

Proof. If $s = 1$ this is the special case of Lemma 5.10 corresponding to the choices $d = c$ and $Q = 0$; in particular, this implies that the exponent t is always at least 1. If $s > 1$ then the result follows by induction using Lemma 5.8(ii). $\square$

For each integer k let $c_\lambda(k) = s$ if k appears in component s of $\mathfrak{t}^\lambda$. Observe that for all k we have $c_\lambda(k) = \min \{ 1 \leq s \leq r \mid k \leq |\lambda^{(1)}| + \dots + |\lambda^{(s)}| \}$. For the next Lemma recall that $\mathfrak{a}_s = |\lambda^{(1)}| + \dots + |\lambda^{(s-1)}|$ and $\mathfrak{b}_s = |\lambda^{(s+1)}| + \dots + |\lambda^{(r)}|$ for $1 \leq s \leq r$.

5.12. Lemma. *Suppose that λ is a multipartition of n . Then*

$$T_{w_\lambda} u_{\lambda'}^- = L_{a_2+1,n}(Q_1) \cdots L_{a_r+1,n}(Q_{r-1}) T_{w_\lambda} + \sum_{k=1}^n \sum_{c=1}^{c_\lambda(k)-1} L_k^c h_{kc}$$

for some $h_{kc} \in H_{k-1,n}$.

Proof. First note that if $r = 1$ then $u_{\lambda'}^- = 1$ and there is nothing to prove; so we assume that $r > 1$. Next, $T_{w_\lambda} u_{\lambda'}^- = T_{w_{\bar{\lambda}}} T_{w_{\lambda/\bar{\lambda}}} u_{\lambda'}^- = T_{w_{\bar{\lambda}}} u_{\bar{\lambda}'}^- T_{w_{\lambda/\bar{\lambda}}}$ by part (iv) of (2.1) since $w_{\lambda/\bar{\lambda}} \in \mathfrak{S}_{\bar{\lambda}'}$. Consequently, it is enough to consider $T_{w_{\bar{\lambda}}} u_{\bar{\lambda}'}^-$; equivalently, we may assume that $\lambda = \bar{\lambda}$ (for clarity we will continue to write $\bar{\lambda}$).

By (5.6), $w_{\bar{\lambda}} = w_{\bar{\lambda},1} \cdots w_{\bar{\lambda},r-1}$ where $w_{\bar{\lambda},s} \in \mathfrak{S}_{\mathfrak{b}_{s-1}}$. Therefore, by (2.1)(iv),

$$\begin{aligned} T_{w_{\bar{\lambda}}} u_{\bar{\lambda}'}^- &= T_{w_{\bar{\lambda},1}} \cdots T_{w_{\bar{\lambda},r-1}} L_{1,\mathfrak{b}_1}(Q_1) \cdots L_{1,\mathfrak{b}_{r-1}}(Q_{r-1}) \\ &= T_{w_{\bar{\lambda},1}} L_{1,\mathfrak{b}_1}(Q_1) \cdots T_{w_{\bar{\lambda},r-1}} L_{1,\mathfrak{b}_{r-1}}(Q_{r-1}). \end{aligned}$$

We also let $n_s = |\lambda^{(s)}|$. Suppose that $1 \leq s \leq r-1$. For $k = 1, \dots, \mathfrak{b}_{s-1}$ set

$$c_{\lambda,s}(k) = \min \{ 0 \leq t \leq r-s \mid k \leq |\lambda^{(s)}| + \cdots + |\lambda^{(s+t)}| \}$$

and let $H^{(s)}$ be the R -submodule of $\mathcal{H}$ spanned by elements of the form $L_k^c h_{kc}$ where $n_s < k \leq \mathfrak{b}_{s-1}$, $1 \leq c \leq c_{\lambda,s}(k)$ and $h_{kc} \in H_{k-1,\mathfrak{b}_{s-1}}$. We claim that

$$T_{w_{\bar{\lambda},s}} L_{1,\mathfrak{b}_s}(Q_s) \cdots T_{w_{\bar{\lambda},r-1}} L_{1,\mathfrak{b}_{r-1}}(Q_{r-1}) \equiv P^{(s)} \pmod{H^{(s)}},$$

where $P^{(s)} = \prod_{t=s}^{r-1} L_{n_s+\cdots+n_t+1,\mathfrak{b}_{s-1}}(Q_t) \cdot T_{w_{\bar{\lambda},s}} \cdots T_{w_{\bar{\lambda},r-1}}$. Taking $s = 1$ will prove the Lemma because $\mathfrak{b}_0 = n$ and $c_\lambda(k) = c_{\lambda,1}(k) + 1$ by the remarks before the Lemma. Note that $P^{(s)}$ has degree $c_{\lambda,s}(k)$ when we consider it as a polynomial in L_k .

To prove the claim we argue by downwards induction on s . If $s \geq c_\lambda(n)$ then $\mathfrak{b}_s = 0$ so that $T_{w_{\bar{\lambda},s}} = L_{1,\mathfrak{b}_s}(Q_s) = P^{(s)} = 1$ and there is nothing to prove. Suppose then that $s < c_\lambda(n)$. Then, by induction,

$$T_{w_{\bar{\lambda},s}} L_{1,\mathfrak{b}_s}(Q_s) \cdots T_{w_{\bar{\lambda},r-1}} L_{1,\mathfrak{b}_{r-1}}(Q_{r-1}) = T_{w_{\bar{\lambda},s}} L_{1,\mathfrak{b}_s}(Q_s) \left\{ P^{(s+1)} + h^{(s+1)} \right\}$$

for some $h^{(s+1)} \in H^{(s+1)}$. Now $\mathfrak{b}_s > 0$ since $s < c_\lambda(n)$, so $\mathfrak{b}_{s-1} = \mathfrak{b}_s + n_s \geq \mathfrak{b}_s \geq 1$. Hence, by Lemma 5.10,

$$\begin{aligned} T_{w_{\bar{\lambda},s}} L_{1,\mathfrak{b}_s}(Q_s) \left\{ P^{(s+1)} + h^{(s+1)} \right\} &= T_{w_{n_s,\mathfrak{b}_s}} L_{1,\mathfrak{b}_s}(Q_s) \left\{ P^{(s+1)} + h^{(s+1)} \right\} \\ &= \left\{ L_{n_s+1,\mathfrak{b}_{s-1}}(Q_s) T_{w_{n_s,\mathfrak{b}_s}} + \sum_{k=n_s+1}^{\mathfrak{b}_{s-1}} L_k h_k T_{w_{n_s,\mathfrak{b}_s}} \right\} \left\{ P^{(s+1)} + h^{(s+1)} \right\}, \end{aligned}$$

for some $h_k \in H_{k-1,k}$. We will move $T_{w_{\bar{\lambda},s}}$ past $P^{(s+1)}$ and $h^{(s+1)}$ using Lemma 5.10.

Viewing $P^{(s+1)}$ as a polynomial in the L_m 's, if $n_{s+1} < m \leq \mathfrak{b}_s$ then the degree of L_m in $P^{(s+1)}$ is at most $c_{\lambda,s+1}(m)$; further, L_m does not appear in $P^{(s+1)}$ if either $m \leq n_{s+1}$ or $m > \mathfrak{b}_s$. Let $p^{(s+1)} = \prod_{t=s+1}^{r-1} L_{n_{s+1}+\cdots+n_t+1,\mathfrak{b}_s}(Q_t)$. Then by

Lemma 5.10 and Corollary 5.11 there exist $h_{k,1}, h_{ck,2} \in H_{k-1,k}$ such that

$$\begin{aligned} T_{w_{n_s, \mathfrak{b}_s}} p^{(s+1)} &= \left\{ L_{n_s+n_{s+1}+1, \mathfrak{b}_{s-1}}(Q_{s+1}) + \sum_{k=n_s+n_{s+1}+1}^{\mathfrak{b}_{s-1}} L_k h_{k,1} \right\} T_{w_{n_s, \mathfrak{b}_s}} p^{(s+2)} \\ &= \left\{ L_{n_s+n_{s+1}+1, \mathfrak{b}_{s-1}}(Q_{s+1}) L_{n_s+n_{s+1}+n_{s+2}+1, \mathfrak{b}_{s-1}}(Q_{s+2}) \right. \\ &\quad \left. + \sum_{k=n_s+n_{s+1}+1}^{\mathfrak{b}_{s-1}} \sum_{c=1}^{c_2(k)} L_k^c h_{ck,2} \right\} T_{w_{n_s, \mathfrak{b}_s}} p^{(s+3)}, \end{aligned}$$

where $c_2(k) = 1$ if $n_s + n_{s+1} < k \leq n_s + n_{s+1} + n_{s+2}$ and $c_2(k) = 2$ if $n_s + n_{s+1} + n_{s+2} < k \leq \mathfrak{b}_{s-1}$. For the second equality we have used Lemma 5.10, Lemma 5.8 and Corollary 5.11. Looking at the definition of $c_{\lambda,s}(m)$, if $n_{s+1} < m \leq \mathfrak{b}_s$ then $c_{\lambda,s+1}(m) = c_{\lambda,s}(m + n_s) - 1$; therefore, continuing in this way we see that

$$T_{w_{n_s, \mathfrak{b}_s}} p^{(s+1)} \equiv \prod_{t=s+1}^{r-1} L_{n_s+n_{s+1}+\dots+n_t+1, \mathfrak{b}_{s-1}}(Q_t) \cdot T_{w_{n_s, \mathfrak{b}_s}} p^{(s)} \pmod{H^{(s)}}.$$

In obtaining this equation notice that if $m > k$ then $L_k h_k L_m h_m = L_m(L_k h_k h_m)$ and $L_k h_k h_m \in H_{m-1,m}$, for any $h_k \in H_{k-1,k}$ and $h_m \in H_{m-1,m}$; similarly, if $m < k$ then $h_k L_m h_m \in H_{k-1,k}$ by Lemma 5.8. It follows that if $L_k^c h$ appears in this expansion, for some $h \in H_{k-1,k}$, then $n_s + n_{s+1} < k \leq \mathfrak{b}_{s-1}$ and so $1 \leq c \leq c_{\lambda,s+1}(k - n_s) = c_{\lambda,s}(k) - 1$ by the remarks above; hence, $L_k h_k L_m h_m \in H^{(s)}$ for all k and m .

Now consider a term from the inductive step of the form $L_k h_k T_{w_{n_s, \mathfrak{b}_s}} p^{(s+1)}$, where $n_s < k \leq \mathfrak{b}_{s-1}$ and $h_k \in H_{k-1,k}$. What we have just shown combined with Lemma 5.8(ii) shows that $L_k h_k T_{w_{n_s, \mathfrak{b}_s}} p^{(s+1)}$ is equal to a linear combination of terms of the form $L_m^c h_{md}$, where $n_s < m \leq \mathfrak{b}_{s-1}$, $1 \leq c \leq c_{\lambda,s}(m)$ and $h_{md} \in H_{m-1,m}$. Moreover, $c \leq c_{\lambda,s}(m)$ with equality only if $k = m > n_s + n_{s+1}$ and $c = 1 = c_{\lambda,s}(m)$ if $n_s < m \leq n_s + n_{s+1}$. Hence, $L_k h_k T_{w_{n_s, \mathfrak{b}_s}} p^{(s+1)} \in H^{(s)}$.

Therefore, combining the last two paragraphs we have shown that

$$T_{w_{n_s, \mathfrak{b}_s}} L_{1, \mathfrak{b}_s}(Q_s) p^{(s+1)} \equiv p^{(s)} \pmod{H^{(s)}}.$$

By similar arguments, the terms $L_{n_s+1, \mathfrak{b}_{s-1}}(Q_s) T_{w_{n_s, \mathfrak{b}_s}} h^{(s+1)}$ and $L_k h_k T_{w_{n_s, \mathfrak{b}_s}} h^{(s+1)}$ from the inductive step also belong to $H^{(s)}$. We leave the details to the reader. $\square$

We are now basically done. The next result essentially computes $\tau(z_\lambda \Psi_{t_\lambda}^*)$; we record it separately because it is also the key to showing that M^λ is a self-dual $\mathcal{H}$ -module and that $\tilde{S}^{\lambda'}$ is isomorphic to the dual of S^λ ; see [24].

5.13. Proposition. *Suppose that λ is a multipartition of n . Then*

$$\tau(z_\lambda T_{w_\lambda}^*) = (-1)^{n(r-1)} q^{\ell(w_\lambda)} \prod_{s=1}^r Q_s^{n-|\lambda^{(s)}|}.$$

Proof. Let $\tilde{u}_{\lambda'}^- = L_{\mathbf{a}_2+1,n}(Q_1) \cdots L_{\mathbf{a}_r+1,n}(Q_{r-1})$. Using (2.3)(i) and Lemma 5.12 shows that

$$\begin{aligned} \tau(z_{\lambda} T_{w_{\lambda}}^*) &= \tau(m_{\lambda} T_{w_{\lambda}} n_{\lambda'} T_{w_{\lambda}}^*) = \tau(x_{\lambda} u_{\lambda}^+ T_{w_{\lambda}} u_{\lambda'}^- y_{\lambda'} T_{w_{\lambda}}^*) \\ &= \tau(u_{\lambda}^+ T_{w_{\lambda}} u_{\lambda'}^- y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}) \\ &= \tau(u_{\lambda}^+ \tilde{u}_{\lambda'}^- T_{w_{\lambda}} y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}) + \tau(u_{\lambda}^+ h y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}), \end{aligned}$$

where $h = \sum_{k=1}^n \sum_{c=1}^{c_{\lambda}(k)-1} L_k^c h_{kc}$ for some $h_{kc} \in H_{k-1,n}$. Consider u_{λ}^+ as a polynomial in L_k . Then the degree of L_k in u_{λ}^+ is $r - c_{\lambda}(k)$. Therefore, if $h_{kc} \neq 0$ then $u_{\lambda}^+ L_k^c h_{kc}$ is a polynomial in L_k which is left divisible by L_k and has degree at most $r - 1$. If $m > k$ then L_m appears in $u_{\lambda}^+ L_k^c h_{kc}$ with exponent at most $r - c_{\lambda}(m) < r$. (If $m < k$ then L_m^d can appear in $u_{\lambda}^+ L_k^c h_{kc}$ for $d \geq r$; however, this does not matter because such terms can be written as a linear combination of Ariki–Koike basis elements in $H_{k-1,k}$.) Therefore, $\tau(u_{\lambda}^+ L_k^c h_{kc} y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}) = 0$ by (2.3)(ii) and, consequently, $\tau(u_{\lambda}^+ h y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}) = 0$; hence,

$$\tau(z_{\lambda} T_{w_{\lambda}}^*) = \tau(u_{\lambda}^+ \tilde{u}_{\lambda'}^- T_{w_{\lambda}} y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}).$$

Considered as polynomials in L_k , u_{λ}^+ has degree $r - c_{\lambda}(k)$ and $\tilde{u}_{\lambda'}^-$ has degree $c_{\lambda}(k) - 1$; consequently, each L_k has degree $r - 1$ in $u_{\lambda}^+ \tilde{u}_{\lambda'}^-$. Therefore, by (2.3) again,

$$\begin{aligned} \tau(z_{\lambda} T_{w_{\lambda}}^*) &= \tau(u_{\lambda}^+ \tilde{u}_{\lambda'}^-) \cdot \tau(T_{w_{\lambda}} y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}) \\ &= \prod_{s=2}^r (-Q_s)^{\mathbf{a}_s} \cdot \prod_{s=1}^{r-1} (-Q_s)^{\mathbf{b}_s} \cdot \tau(T_{w_{\lambda}} y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}) \\ &= (-1)^{n(r-1)} \prod_{s=1}^r Q_s^{n-|\lambda^{(s)}|} \cdot \tau(T_{w_{\lambda}} y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}), \end{aligned}$$

since $\mathbf{a}_s + \mathbf{b}_s = n - |\lambda^{(s)}|$ for all s (and $\mathbf{a}_1 = 0 = \mathbf{b}_r$). To complete the proof recall that $\mathfrak{S}_{\lambda} \cap {}^{w_{\lambda}}\mathfrak{S}_{\lambda'} = \{1\}$ and that w_{λ} is a distinguished $(\mathfrak{S}_{\lambda}, \mathfrak{S}_{\lambda'})$ -double coset representative (consider the tableaux $\mathbf{t}^{\lambda}$ and $\mathbf{t}_{\lambda}$), so

$$T_{w_{\lambda}} y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda} = \sum_{\substack{u \in \mathfrak{S}_{\lambda'} \\ v \in \mathfrak{S}_{\lambda}}} (-q)^{-\ell(u)} T_{w_{\lambda}} T_u T_{w_{\lambda}}^* T_v = \sum_{\substack{u \in \mathfrak{S}_{\lambda'} \\ v \in \mathfrak{S}_{\lambda}}} (-q)^{-\ell(u)} T_{w_{\lambda}} T_{uw_{\lambda}^{-1}v}.$$

Therefore, $\tau(T_{w_{\lambda}} y_{\lambda'} T_{w_{\lambda}}^* x_{\lambda}) = q^{\ell(w_{\lambda})}$ by (2.3)(ii) (corresponding to $u = v = 1$), so the Proposition follows. $\square$

We can now give our first formula for the Schur elements $s_{\lambda}(\mathbf{q})$ of $\mathcal{H}$.

5.14. Corollary. *Suppose that λ is a multipartition of n . Then*

$$s_{\lambda}(\mathbf{q}) = (-1)^{n(r-1)} q^{-\ell(w_{\lambda})} \gamma_{\mathbf{t}_{\lambda}} \gamma'_{\mathbf{t}_{\lambda'}} \prod_{s=1}^r Q_s^{|\lambda^{(s)}| - n}.$$

Proof. By Lemma 2.6, $s_{\lambda}(\mathbf{q}) = 1/\tau(F_{\mathbf{t}_{\lambda}})$ and $F_{\mathbf{t}_{\lambda}} = \frac{1}{\gamma_{\mathbf{t}_{\lambda}} \gamma'_{\mathbf{t}_{\lambda'}}} z_{\lambda} \Psi_{\mathbf{t}_{\lambda}}^*$ by Corollary 5.2; therefore, $s_{\lambda}(\mathbf{q}) = \gamma_{\mathbf{t}_{\lambda}} \gamma'_{\mathbf{t}_{\lambda'}} / \tau(z_{\lambda} \Psi_{\mathbf{t}_{\lambda}}^*)$. Now, τ is a trace form by (2.3)(i), so

$$\tau(z_{\lambda} \Psi_{\mathbf{t}_{\lambda}}^*) = \tau(x_{\lambda} u_{\lambda}^+ T_{w_{\lambda}} u_{\lambda'}^- y_{\lambda'} \Psi_{\mathbf{t}_{\lambda}}^*) = \tau(u_{\lambda}^+ T_{w_{\lambda}} u_{\lambda'}^- y_{\lambda'} \Psi_{\mathbf{t}_{\lambda}}^* x_{\lambda}).$$

It is well-known and easy to check that the permutation w_λ has the “trivial intersection property”; that is, $\mathfrak{S}_\lambda \cap {}^w\mathfrak{S}_{\lambda'} \neq \{1\}$ if and only if $\mathfrak{S}_\lambda w_\lambda \mathfrak{S}_{\lambda'} = \mathfrak{S}_\lambda w_\lambda \mathfrak{S}_{\lambda'}$. Therefore, $y_{\lambda'} T_w^* x_\lambda \neq 0$ if and only if $w \in \mathfrak{S}_\lambda w_\lambda \mathfrak{S}_{\lambda'}$ (see, for example, [9, (4.9)]). Now, $\Psi_{t_\lambda}^* = T_{w_\lambda}^* + \sum_{w < w_\lambda} p_{t_\lambda w} T_w^*$ by Proposition 5.1(ii); so $y_{\lambda'} \Psi_{t_\lambda}^* x_\lambda = y_{\lambda'} T_{w_\lambda}^* x_\lambda$ since w_λ is the unique element of minimal length in $\mathfrak{S}_\lambda w_\lambda \mathfrak{S}_{\lambda'}$. Therefore,

$$\tau(z_\lambda \Psi_{t_\lambda}^*) = \tau(u_\lambda^+ T_{w_\lambda} u_{\lambda'}^- y_{\lambda'} T_{w_\lambda}^* x_{\lambda'}) = \tau(x_\lambda u_\lambda^+ T_{w_\lambda} u_{\lambda'}^- y_{\lambda'} T_{w_\lambda}^*) = \tau(z_\lambda T_{w_\lambda}^*).$$

The result now follows from Proposition 5.13. $\square$

A closed formula for $\gamma'_{t_{\lambda'}}$ is given by (3.7)(i); therefore, in order to find an explicit formula for $s_\lambda(\mathbf{q})$ we need only compute γ_{t_λ} . Although the formula below looks formidable, its proof follows readily enough from the definition of γ_{t_λ} .

Recall that the ij th hook in the diagram $[\lambda^{(s)}]$ is the collection of nodes to the right of and below the node (i, j, s) , including the node (i, j, s) itself. The ij th hook length $h_{ij}^{\lambda^{(s)}} = \lambda_i^{(s)} + \lambda_j^{(s)'} - i - j + 1$ is the number of nodes in the ij th hook and the leg length, $\ell_{ij}^{\lambda^{(s)}} = \lambda_j^{(s)'} - j + 1$, is the number of nodes in the “leg” of this hook. Observe that if (a, b, c) and (i, j, c) are two removable nodes in $[\lambda^{(c)}]$ with $a \leq i$ and $j \leq b$ then $h_{aj}^{\lambda^{(c)}} = b - a - j + i + 1$.

5.15. Lemma. *Suppose that λ is a multipartition of n . Then*

$$\gamma_{t_\lambda} = q^{\ell(w_\lambda)} \prod_{(i,j,s) \in [\lambda]} \frac{[h_{ij}^{\lambda^{(s)}}]_q}{[\ell_{ij}^{\lambda^{(s)}}]_q} \prod_{t=s+1}^r (q^{j-i} Q_s - q^{\lambda_1^{(t)}} Q_t) \prod_{k=1}^{\lambda_1^{(t)}} \frac{(q^{j-i} Q_s - q^{k-1-\lambda_k^{(t)'}} Q_t)}{(q^{j-i} Q_s - q^{k-\lambda_k^{(t)'}} Q_t)}.$$

(Note that $\lambda_k^{(t)'}$ is the length of the k^{th} column of $[\lambda^{(t)}]$.)

Proof. We argue by induction on n . If $n = 0$, both sides are 1 and there is nothing to prove (by convention, empty products are 1). Suppose that $n > 0$. Let $\mu = \text{Shape}(t_\lambda \downarrow (n-1))$; then μ is a multipartition of $n-1$. Recall that $\alpha(\lambda) = \frac{1}{2} \sum_{s=1}^r \sum_{i \geq 1} (\lambda_i^{(s)} - 1) \lambda_i^{(s)}$. Applying the definitions (see (3.6)),

$$\frac{\gamma_{t_\lambda}}{\gamma_{t_\mu}} = q^{\ell(w_\lambda) + \alpha(\lambda) - \ell(w_\mu) - \alpha(\mu)} \frac{\prod_{x \in \mathcal{A}_{t_\lambda}(n)} (\text{res}_{t_\lambda}(n) - \text{res}(x))}{\prod_{y \in \mathcal{R}_{t_\lambda}(n)} (\text{res}_{t_\lambda}(n) - \text{res}(y))}.$$

Assume that n appears in row a and column b of $t_\lambda^{(c)}$. First consider the contribution that the addable and removable nodes in $[\lambda^{(c)}]$ make to γ_{t_λ} . Looking at the definitions above (3.6), these nodes occur in pairs (x, y) where $y < (a, b, c)$ is a removable node in row i and x is an addable node in row $i+1$ for some $i \geq a$. If x is in column d of $[\lambda^{(c)}]$ and y is in column d' then $d \leq d' < b$ and

$$\begin{aligned} \frac{\text{res}_{t_\lambda}(n) - \text{res}_{t_\lambda}(x)}{\text{res}_{t_\lambda}(n) - \text{res}_{t_\lambda}(y)} &= \frac{q^{b-a} Q_c - q^{d-(i+1)} Q_c}{q^{b-a} Q_c - q^{d'-i} Q_c} = \prod_{j=d}^{d'} \frac{q^{b-a} - q^{j-(i+1)}}{q^{b-a} - q^{j-i}} \\ &= \prod_{j=d}^{d'} \frac{q^{j-i-1} (q^{b-a-j+i+1} - 1)}{q^{j-i} (q^{b-a-j+i} - 1)} = \prod_{j=d}^{d'} \frac{q^{-1} (q^{h_{aj}^{\lambda^{(c)}}} - 1)}{(q^{h_{aj}^{\lambda^{(c)}}} - 1)} \\ &= \prod_{j=d}^{d'} \frac{q^{-1} [h_{aj}^{\lambda^{(c)}}]_q}{[h_{aj}^{\lambda^{(c)}} - 1]_q}. \end{aligned}$$

Therefore,

$$\frac{\prod_{(i,j,c) \in \mathcal{A}_{t_\lambda}(n)} (\text{res}_{t_\lambda}(n) - \text{res}_{t_\lambda}(i, j, c))}{\prod_{(i,j,c) \in \mathcal{R}_{t_\lambda}(n)} (\text{res}_{t_\lambda}(n) - \text{res}_{t_\lambda}(i, j, c))} = \prod_{j=1}^{b-1} \frac{q^{-1} [h_{aj}^{\lambda^{(c)}}]_q}{[h_{aj}^{\lambda^{(c)}} - 1]_q} = q^{1-b} \prod_{j=1}^{b-1} \frac{[h_{aj}^{\lambda^{(c)}}]_q}{[h_{aj}^{\mu^{(c)}}]_q}.$$

Note that $\alpha(\lambda) = \alpha(\mu) + b - 1$; hence, by induction, this accounts for the factor $q^{\ell(w_\lambda)}$ in Lemma 5.15.

Now γ_{t_μ} is known by induction and it contains as a factor the left hand term in the product below. Further, $\ell_{ij}^{\mu^{(s)}} = \ell_{ij}^{\lambda^{(s)}}$ and $h_{ij}^{\mu^{(s)}} = h_{ij}^{\lambda^{(s)}}$ if $(i, s) \neq (a, c)$, and $\ell_{aj}^{\mu^{(c)}} = \ell_{aj}^{\lambda^{(c)}}$ for $1 \leq j < b$, so

$$\begin{aligned} \left(\prod_{(i,j,s) \in [\mu]} \frac{[h_{ij}^{\mu^{(s)}}]_q}{[\ell_{ij}^{\mu^{(s)}}]_q} \right) \left(\prod_{j=1}^{b-1} \frac{[h_{aj}^{\lambda^{(c)}}]_q}{[h_{aj}^{\mu^{(c)}}]_q} \right) &= \left(\prod_{\substack{(i,j,s) \in [\lambda] \\ (i,s) \neq (a,c)}} \frac{[h_{ij}^{\lambda^{(s)}}]_q}{[\ell_{ij}^{\lambda^{(s)}}]_q} \right) \left(\prod_{j=1}^{b-1} \frac{[h_{aj}^{\lambda^{(c)}}]_q}{[\ell_{aj}^{\lambda^{(c)}}]_q} \right) \\ &= \prod_{(i,j,s) \in [\lambda]} \frac{[h_{ij}^{\lambda^{(s)}}]_q}{[\ell_{ij}^{\lambda^{(s)}}]_q}, \end{aligned}$$

since $[h_{ab}^{\lambda^{(c)}}]_q = 1 = [\ell_{ab}^{\lambda^{(c)}}]_q$. This accounts for the left hand factor in the expression for γ_{t_λ} given in the statement of the Lemma.

Finally, consider the nodes in $\mathcal{A}_{t_\lambda}(n)$ and $\mathcal{R}_{t_\lambda}(n)$ which are in component t for some $t > c$ (there are no such nodes for $t < c$). Again, almost all of the addable and removable nodes in component t occur in pairs placed in consecutive rows; however, this time there is also an additional addable node at the end of the first row of $\lambda^{(t)}$. As above, it is easier to insert extra factors which cancel out and so take a product over all of the columns of $\lambda^{(t)}$. An argument similar to that above shows that the nodes in $\mathcal{A}_{t_\lambda}(n)$ and $\mathcal{R}_{t_\lambda}(n)$ which do not belong to component c contribute the factor

$$\prod_{t=c+1}^r (q^{b-a} Q_c - q^{\lambda_1^{(t)}} Q_t) \prod_{k=1}^{\lambda_1^{(t)}} \frac{(q^{b-a} Q_c - q^{k-1-\lambda_k^{(t)'}} Q_t)}{(q^{b-a} Q_c - q^{k-\lambda_k^{(t)'}} Q_t)}$$

to γ_{t_λ} . Using induction to combine the formulae above proves the Lemma. $\square$

We can now give a closed formula for the Schur elements.

5.16. Corollary. *Suppose that λ is a multipartition of n and for $1 \leq s < t \leq r$ let*

$$X_{st}^\lambda = \prod_{(i,j) \in [\lambda^{(t)}]} (q^{j-i} Q_t - Q_s) \cdot \prod_{(i,j) \in [\lambda^{(s)}]} (q^{j-i} Q_s - q^{\lambda_1^{(t)}}) \prod_{k=1}^{\lambda_1^{(t)}} \frac{(q^{j-i} Q_s - q^{k-1-\lambda_k^{(t)'}} Q_t)}{(q^{j-i} Q_s - q^{k-\lambda_k^{(t)'}} Q_t)}.$$

Then

$$s_\lambda(\mathbf{q}) = (-1)^{n(r-1)} (Q_1 \dots Q_r)^{-n} q^{-\alpha(\lambda')} \prod_{s=1}^r \prod_{(i,j) \in [\lambda^{(s)}]} Q_s [h_{ij}^{\lambda^{(s)}}]_q \cdot \prod_{1 \leq s < t \leq r} X_{st}^\lambda.$$

Proof. Looking at the definitions, $[k]_q' = 1 + q^{-1} + \dots + q^{1-k} = q^{1-k} [k]_q$; therefore, $([k]_q^!)^{\prime} = q^{-\frac{1}{2}k(k-1)} [k]_q^!$; consequently, $([\lambda']_q^!)^{\prime} = q^{-\alpha(\lambda')} [\lambda']_q^!$. Next observe that $[\lambda']_q^! = \prod_{(i,j,s) \in [\lambda]} [\ell_{ij}^{\lambda^{(s)}}]_q$. Further, $(i, j) \in [\lambda^{(s)}]$ if and only if $(j, i) \in [\lambda'^{(r-s+1)}]$.

Therefore, applying ' to (3.7)(i) (and swapping the roles of s and t in the right hand factor),

$$\gamma'_{t\lambda'} = q^{-\alpha(\lambda')} \left(\prod_{(i,j,s) \in [\lambda]} [\ell_{ij}^{\lambda^{(s)}}]_q \right) \left(\prod_{1 \leq s < t \leq r} \prod_{(i,j) \in [\lambda^{(t)}]} (q^{j-i} Q_t - Q_s) \right).$$

By Corollary 5.14, $s_\lambda(\mathbf{q}) = (-1)^{n(r-1)} q^{-\ell(w_\lambda)} \gamma_{t\lambda} \gamma'_{t\lambda'} \prod_{s=1}^r Q_s^{|\lambda^{(s)}| - n}$; so the result now follows by Lemma 5.15. $\square$

5.17. *Example.* It is straightforward to check that Corollary 5.16 gives the same rational functions for the Schur elements $s_{\eta_t}(\mathbf{q})$ as were obtained in Example 2.7.

It is not at all obvious that the formula in Corollary 5.16 for the Schur elements agrees with that conjectured by Malle, so we now show that this is the case. This takes quite a lot of work; however, it also results in a more symmetrical formula for $s_\lambda(\mathbf{q})$. To do this we need to rewrite the Schur elements as functions of beta numbers (or first column hook lengths).

Define the length of a partition σ to be the smallest integer $\ell(\sigma)$ such that $\sigma_i = 0$ for all $i > \ell(\sigma)$. The length of a multipartition λ is $\ell(\lambda) = \max \{ \ell(\lambda^{(s)}) \mid 1 \leq s \leq r \}$.

Fix an integer L such that $L \geq \ell(\lambda)$. The L -beta numbers for $\lambda^{(s)}$ are the integers $\beta_i^{(s)} = \lambda_i^{(s)} + L - i$ for $i = 1, \dots, L$; note that $\beta_1^{(s)} > \dots > \beta_L^{(s)} \geq 0$. Malle calls the $r \times L$ matrix $B = (\beta_i^{(s)})_{s,i}$ the L -symbol of λ . Actually, this is not quite one of Malle's symbols; it is what Broué and Kim [6] call B an *ordinary symbol*. We also let $B_s = \{\beta_1^{(s)}, \dots, \beta_L^{(s)}\}$ for $s = 1, \dots, r$.

If we change L to $L + 1$ the beta set B_s is *shifted* to $\{\beta_1^{(s)} + 1, \dots, \beta_L^{(s)} + 1, 0\}$. A function of beta numbers is *invariant under beta shifts* if it is unchanged by such transformations; equivalently, the function is independent of L provided that L is large enough. For example, the formula for $s_\lambda(\mathbf{q})$ below is invariant under beta shifts since $s_\lambda(\mathbf{q})$ does not depend on L .

5.18. Theorem. *Suppose that λ is a multipartition of n with L -symbol $B = (\beta_i^{(s)})_{s,i}$ such that $L \geq \ell(\lambda)$. Then*

$$s_\lambda(\mathbf{q}) = (-1)^{a_{rL}} q^{b_{rL}} \frac{\prod_{1 \leq s < t \leq r} (Q_s - Q_t)^L \cdot \prod_{1 \leq s, t \leq r} \prod_{\alpha_s \in B_s} \prod_{1 \leq k \leq \alpha_s} (q^k Q_s - Q_t)}{(q-1)^n (Q_1 \dots Q_r)^n \prod_{1 \leq s \leq t \leq r} \prod_{\substack{(\alpha_s, \alpha_t) \in B_s \times B_t \\ \alpha_s > \alpha_t \text{ if } s=t}} (q^{\alpha_s} Q_s - q^{\alpha_t} Q_t)},$$

where $a_{rL} = n(r-1) + \binom{r}{2} \binom{L}{2}$ and $b_{rL} = \frac{rL(L-1)(2rL-r-3)}{12}$.

Proof. Adopting the notation of Corollary 5.16,

$$s_\lambda(\mathbf{q}) = (-1)^{n(r-1)} (Q_1 \dots Q_r)^{-n} q^{-\alpha(\lambda')} \prod_{s=1}^r \prod_{(i,j) \in [\lambda^{(s)}]} Q_s [h_{ij}^{\lambda^{(s)}}]_q \cdot \prod_{1 \leq s < t \leq r} X_{st}^\lambda.$$

We consider each of these factors separately.

First, let σ be a partition of m with beta numbers $(\beta_1, \dots, \beta_L)$, where $L \geq \ell(\sigma)$. Then

$$\prod_{(i,j) \in [\sigma]} [h_{ij}^\sigma]_q = (q-1)^{-m} \prod_{(i,j) \in [\sigma]} (q^{h_{ij}^\sigma} - 1) = (q-1)^{-m} \frac{\prod_{i=1}^L \prod_{k=1}^{\beta_i} (q^k - 1)}{\prod_{1 \leq i < j \leq L} (q^{\beta_i - \beta_j} - 1)}.$$

This is quite well-known and is easily proved by first observing that the right hand side is invariant under beta shifts and then by arguing by induction on the number of columns.

Ignoring the leading term $(q-1)^m$ on the right hand side, the number of factors in the numerator is $m + \binom{L}{2}$, whereas the number of factors in the denominator is $\binom{L}{2}$. Consequently, multiplying the left hand side by Q^m , say, is the same as multiplying each factor, top and bottom, on the right hand side by Q . Therefore, returning to the multipartition λ ,

$$q^{-\alpha(\lambda')} \prod_{s=1}^r \prod_{(i,j) \in [\lambda^{(s)}]} Q_s [h_{ij}^{\lambda^{(s)}}]_q = q^{N-\alpha(\lambda')} (q-1)^{-n} \prod_{s=1}^r \frac{\prod_{i=1}^L \prod_{k=1}^{\beta_i^{(s)}} (q^k Q_s - Q_s)}{\prod_{1 \leq i < j \leq L} (q^{\beta_i^{(s)}} Q_s - q^{\beta_j^{(s)}} Q_s)},$$

where $N = \sum_{s=1}^r \sum_{j=1}^L (j-1)\beta_j^{(s)}$. It is well-known (in the right circles) and easy enough to check that $\alpha(\lambda') = \sum_{s=1}^r \sum_{j=1}^L (j-1)\lambda_j^{(s)}$; therefore, $N - \alpha(\lambda') = r \binom{L}{3}$.

Fix s and t with $1 \leq s < t \leq r$; there are $\binom{r}{2}$ such choices. A quick calculation shows that $b_{rL} = r \binom{L}{3} + \binom{r}{2} \sigma(L)$, where $\sigma(L) = (2L-1)L(L-1)/6 = \sum_{j=1}^{L-1} j^2$. Therefore, in order to complete the proof it is enough to show that $X_{st}^\lambda = Y_{st}^\lambda$ where

$$Y_{st}^\lambda = (-1)^{\binom{L}{2}} q^{\sigma(L)} \frac{\prod_{\alpha_s \in B_s} \prod_{1 \leq k \leq \alpha_s} (q^k Q_s - Q_t) \cdot \prod_{\alpha_t \in B_t} \prod_{1 \leq k \leq \alpha_t} (q^k Q_t - Q_s)}{\prod_{(\alpha_s, \alpha_t) \in B_s \times B_t} (q^{\alpha_s} Q_s - q^{\alpha_t} Q_t)},$$

It is not hard to see that Y_{st}^λ is invariant under beta shifts so we may change L arbitrarily, provided that $L \geq \max\{\ell(\lambda^{(s)}), \ell(\lambda^{(t)})\}$. We prove our claim that $X_{st}^\lambda = Y_{st}^\lambda$ by induction in three incremental steps. We start the induction by observing that both products are equal to 1 when $\lambda^{(s)} = \lambda^{(t)} = (0)$ — to see this it is easiest to take $L = 0$.

Next consider the case where $\lambda^{(t)} = (0)$. Since Y_{st}^λ is invariant under beta shifts we may assume that $\ell(\lambda^{(s)}) = L$. Assume by way of induction that we have proved the claim for $(\lambda_1^{(s)}, \dots, \lambda_{L-1}^{(s)})$. Adding a non-empty L^{th} row to $\lambda^{(s)}$ changes X_{st}^λ

by the factor

$$\begin{aligned}
\prod_{k=1-L}^{\lambda_L^{(s)}-L} (q^k Q_s - Q_t) &= \prod_{k=1}^{L-1} q^{-k} (Q_s - q^k Q_t) \cdot (Q_s - Q_t) \cdot \prod_{k=1}^{\lambda_L^{(s)}-L} (q^k Q_s - Q_t) \\
&= (-1)^{L-1} \frac{\prod_{1 \leq k \leq L-1} (q^k Q_t - Q_s) \cdot (Q_s - Q_t) \cdot \prod_{1 \leq k \leq \lambda_L^{(s)}} (q^k Q_s - Q_t)}{q^{\binom{L}{2}} \prod_{\lambda_L^{(s)}-L+1 \leq k \leq \lambda_L^{(s)}} (q^k Q_s - Q_t)} \\
&= (-1)^{L-1} \frac{\prod_{1 \leq k \leq \beta_L^{(t)}} (q^k Q_t - Q_s) \cdot (Q_s - Q_t) \cdot \prod_{1 \leq k \leq \beta_L^{(s)}} (q^k Q_s - Q_t)}{\prod_{\alpha_t \in B_t} (q^{\beta_L^{(s)}} Q_s - q^{\alpha_t} Q_t)},
\end{aligned}$$

the last equality following because $\beta_L^{(s)} = \lambda_L^{(s)}$ and $\beta_i^{(t)} = L - i$ for $i = 1, \dots, L$. Notice that the $(L-1)$ -beta numbers for $(\lambda_1^{(s)}, \dots, \lambda_{L-1}^{(s)})$ all increase by 1 when we add the extra row $\lambda_L^{(s)}$ to $\lambda^{(s)}$. Let B'_s be the set of $(L-1)$ -beta numbers for $(\lambda_1^{(s)}, \dots, \lambda_{L-1}^{(s)})$ and let $B'_t = \{0, 1, \dots, L-2\}$ be the set of $(L-1)$ -beta numbers for $\lambda^{(t)}$. Then

$$\begin{aligned}
\prod_{(\alpha_s, \alpha_t) \in B_s \times B_t} (q^{\alpha_s} Q_s - q^{\alpha_t} Q_t) &= \prod_{1 \leq i \leq L} \prod_{0 \leq k < L} (q^{\beta_i^{(s)}} Q_s - q^k Q_t) \\
&= \prod_{\alpha_t \in B_t} (q^{\beta_L^{(s)}} Q_s - q^{\alpha_t} Q_t) \cdot \prod_{1 \leq i < L} (q^{\beta_i^{(s)}} Q_s - Q_t) \cdot \prod_{(\alpha_s, \alpha_t) \in B'_s \times B'_t} q (q^{\alpha_s} Q_s - q^{\alpha_t} Q_t).
\end{aligned}$$

This equation allows us to rewrite the denominator of the preceding equation and so see that the change in X_{st}^λ is the same as the change in Y_{st}^λ (in particular, the change of the scalar is $(-1)^{L-1} q^{(L-1)^2}$ in both cases). This proves our claim when $\lambda^{(t)} = (0)$.

The next step is to fix $\lambda^{(s)}$ and assume that $\lambda^{(t)} = (a)$ for some $a \geq 0$. If $\lambda^{(s)} = (0)$ in this case then it is straightforward to check the claim (or to modify the argument below), so assume that $\lambda^{(s)} \neq (0)$ and let $L = \ell(\lambda^{(s)}) \geq 1 \geq \ell(\lambda^{(t)})$. The case $a = 0$ we already understand. Next, changing $\lambda^{(t)}$ from $(a-1)$ to (a) changes X_{st}^λ by the factor

$$\begin{aligned}
&(q^{a-1} Q_t - Q_s) \prod_{(i,j) \in [\lambda^{(s)}]} \frac{(q^{j-i} Q_s - q^a Q_t)}{(q^{j-i} Q_s - q^{a-1} Q_t)} \frac{(q^{j-i} Q_s - q^{a-2} Q_t)}{(q^{j-i} Q_s - q^{a-1} Q_t)} \\
&= (q^{a-1} Q_t - Q_s) \prod_{(i,j) \in [\lambda^{(s)}]} \frac{(q^{a-1} Q_t - q^{j-i-1} Q_s)(q^{j-i+1} Q_s - q^{a-1} Q_t)}{(q^{j-i} Q_s - q^{a-1} Q_t)(q^{a-1} Q_t - q^{j-i} Q_s)} \\
&= (q^{a-1} Q_t - Q_s) \prod_{i=1}^L \frac{(q^{a-1} Q_t - q^{-i} Q_s)}{(q^{a-1} Q_t - q^{1-i} Q_s)} \frac{(q^{\lambda_i^{(s)}-i+1} Q_s - q^{a-1} Q_t)}{(q^{\lambda_i^{(s)}-i} Q_s - q^{a-1} Q_t)} \\
&= (q^{a-1} Q_t - q^{-L} Q_s) \prod_{i=1}^L \frac{(q^{\beta_i^{(s)}+1} Q_s - q^{\beta_1^{(t)}} Q_t)}{(q^{\beta_i^{(s)}} Q_s - q^{\beta_1^{(t)}} Q_t)} \\
&= (q^{\beta_1^{(t)}} Q_t - Q_s) \prod_{i=1}^L \frac{(q^{\beta_i^{(s)}} Q_s - q^{\beta_1^{(t)}-1} Q_t)}{(q^{\beta_i^{(s)}} Q_s - q^{\beta_1^{(t)}} Q_t)}.
\end{aligned}$$

This last product is exactly the change in Y_{st}^λ so we now know that $X_{st}^\lambda = Y_{st}^\lambda$ when $\lambda^{(t)} = (a)$.

Finally, suppose that $\lambda^{(t)}$ has more than one row. For convenience, let $l = \ell(\lambda^{(s)})$, $m = \ell(\lambda^{(t)}) > 1$ and $b = \lambda_m^{(t)}$ and assume that $L \geq l, m$. When we add row m to $\lambda^{(t)}$, X_{st}^λ changes by the factor

$$\begin{aligned}
& \prod_{j=1}^b (q^{j-m} Q_t - Q_s) \cdot \prod_{(i,j) \in [\lambda^{(s)}]} \prod_{k=1}^b \frac{(q^{j-i} Q_s - q^{k-m-1} Q_t)}{(q^{j-i} Q_s - q^{k-m} Q_t)} \frac{(q^{j-i} Q_s - q^{k-m+1} Q_t)}{(q^{j-i} Q_s - q^{k-m} Q_t)} \\
&= \prod_{j=1}^b (q^{j-m} Q_t - Q_s) \cdot \prod_{(i,j) \in [\lambda^{(s)}]} \prod_{k=1}^b \frac{(q^{j-i+1} Q_s - q^{k-m} Q_t)}{(q^{j-i} Q_s - q^{k-m} Q_t)} \frac{(q^{k-m} Q_t - q^{j-i-1} Q_s)}{(q^{k-m} Q_t - q^{j-i} Q_s)} \\
&= \prod_{j=1}^b (q^{j-m} Q_t - Q_s) \cdot \prod_{i=1}^l \prod_{k=1}^b \frac{q(q^{\lambda_i^{(s)}-i} Q_s - q^{k-m-1} Q_t)}{(q^{\lambda_i^{(s)}-i} Q_s - q^{k-m} Q_t)} \frac{(q^{k-m} Q_t - q^{j-i} Q_s)}{(q^{k-m} Q_t - q^{j-i-1} Q_s)} \\
&= \prod_{j=1}^b (q^{j-m} Q_t - Q_s) \cdot q^{lb} \prod_{i=1}^l \frac{(q^{\lambda_i^{(s)}-i} Q_s - q^{k-m} Q_t)}{(q^{\lambda_i^{(s)}-i} Q_s - q^{b-m} Q_t)} \cdot \prod_{k=1}^b \frac{(q^{k-m} Q_t - q^{j-l} Q_s)}{(q^{k-m} Q_t - Q_s)} \\
&= \prod_{k=1}^b (q^{k-m+l} Q_t - Q_s) \cdot \prod_{i=1}^l \frac{(q^{\beta_i^{(s)}} Q_s - q^{\beta_m^{(t)}-b} Q_t)}{(q^{\beta_i^{(s)}} Q_s - q^{\beta_m^{(t)}} Q_t)} \\
&= \prod_{k=L-m+1}^{\beta_m^{(t)}} (q^k Q_t - Q_s) \cdot \prod_{i=1}^L \frac{(q^{\beta_i^{(s)}} Q_s - q^{\beta_m^{(t)}-b} Q_t)}{(q^{\beta_i^{(s)}} Q_s - q^{\beta_m^{(t)}} Q_t)},
\end{aligned}$$

where the last line follows by a short calculation since $\beta_m^{(t)} = b + L - m$ (and $l \leq L$). As $\beta_m^{(t)}$ is the only beta number that has changed, this factor is precisely the change in Y_{st}^λ when an extra row is added to $\lambda^{(t)}$.

We have now shown that $X_{st}^\lambda = Y_{st}^\lambda$ in all cases, so the theorem is proved. $\square$

Finally, comparing Theorem 5.18 with [6, Prop 3.17] we see that our formula for the Schur elements agrees with Malle's — as it must because Malle's conjecture has already been proved by Geck, Iancu and Malle [15]. Actually, there is still a small amount of work to be done in reconciling the two formulas because Broué and Kim [6] write the exponent of q as a sum of binomial coefficients; however, their expression simplifies to give b_{rL} .

Acknowledgements

I would like to thank the referee for many useful comments and corrections and Jean Michel for writing some CHEVIE code for computing the Schur elements.

REFERENCES

- [1] S. ARIKI, *On the semi-simplicity of the Hecke algebra of $(\mathbb{Z}/r\mathbb{Z}) \wr \mathfrak{S}_n$* , J. Algebra, **169** (1994), 216–225.
- [2] ———, *Representation theory of a Hecke algebra of $G(r, p, n)$* , J. Algebra, **177** (1995), 164–185.
- [3] ———, *On the decomposition numbers of the Hecke algebra of $G(m, 1, n)$* , J. Math. Kyoto Univ., **36** (1996), 789–808.
- [4] S. ARIKI AND K. KOIKE, *A Hecke algebra of $(\mathbb{Z}/r\mathbb{Z}) \wr \mathfrak{S}_n$ and construction of its irreducible representations*, Adv. Math., **106** (1994), 216–243.
- [5] K. BREMKKE AND G. MALLE, *Reduced words and a length function for $G(e, 1, n)$* , Indag. Math., **8** (1997), 453–469.

- [6] M. BROUÉ AND S. KIM, *Sur les familles de caractères des algèbres de hecke cyclotomiques*, preprint 2001.
- [7] M. BROUÉ AND G. MALLE, *Zyklotomische Heckealgebren*, *Asterisque*, **212** (1993), 119–189.
- [8] R. W. CARTER, *Finite Groups of Lie Type*, John Wiley, New York, 1985.
- [9] R. DIPPER AND G. JAMES, *Representations of Hecke algebras of general linear groups*, *Proc. L.M.S.* (3), **52** (1986), 20–52.
- [10] ———, *Blocks and idempotents of Hecke algebras of general linear groups*, *Proc. L.M.S.* (3), **54** (1987), 57–82.
- [11] ———, *Representations of Hecke algebras of type B_n* , *J. Algebra*, **146** (1992), 454–481.
- [12] R. DIPPER, G. JAMES, AND A. MATHAS, *Cyclotomic q -Schur algebras*, *Math. Z.*, **229** (1999), 385–416.
- [13] J. DU AND H. RUI, *Specht modules for Ariki-Koike algebras*, *Comm. Alg.*, **29** (2001), 4710–4719.
- [14] J. S. FRAME, G. DE. B. ROBINSON, AND R. M. THRALL, *The hook graphs of the symmetric group*, *Canad. J. Math.*, **6** (1954), 316–324.
- [15] M. GECK, L. IANCU, AND G. MALLE, *Weights of Markov traces and generic degrees*, *Indag. Math.*, **11** (2000), 379–397.
- [16] P. N. HOEFSMIT, *Representations of Hecke algebras of finite groups with BN -pairs of classical type*, PhD thesis, University of British Columbia, 1979.
- [17] G. D. JAMES AND A. MATHAS, *The Jantzen sum formula for cyclotomic q -Schur algebras*, *Trans. A.M.S.*, **352** (2000), 5381–5404.
- [18] G. LUSZTIG, *On a theorem of Benson and Curtis*, *J. Algebra*, **71** (1981), 490–498.
- [19] G. MALLE, *Unipotente Grade imprimitiver komplexer Spiegelungsgruppen*, *J. Algebra*, **177** (1995), 768–826.
- [20] ———, *On the generic degrees of cyclotomic algebras*, *Represent. Theory*, **4** (2000), 342–369 (electronic).
- [21] G. MALLE AND A. MATHAS, *Symmetric cyclotomic Hecke algebras*, *J. Algebra*, **205** (1998), 275–293.
- [22] A. MATHAS, *Hecke algebras and Schur algebras of the symmetric group*, *Univ. Lecture Notes*, **15**, A.M.S., 1999.
- [23] ———, *Murphy operators and the centre of the Iwahori-Hecke algebras of type A* , *J. Alg. Comb.*, **9** (1999), 295–313.
- [24] ———, *Tilting modules for cyclotomic Schur algebras*, in preparation 2001.
- [25] G. E. MURPHY, *On the representation theory of the symmetric groups and associated Hecke algebras*, *J. Algebra*, **152** (1992), 492–513.

SCHOOL OF MATHEMATICS AND STATISTICS F07
 UNIVERSITY OF SYDNEY, SYDNEY N.S.W. 2006. mathas@maths.usyd.edu.au
 AUSTRALIA. www.maths.usyd.edu.au/u/mathas/